

Baseline drug-resistance mutations are detectable in HCV genes NS3 and NS5A but not NS5B in acute and chronic HIV-coinfected patients

Adele L. McCormick^{1*}

Luke Moynihan¹

Malcolm J Macartney¹

Ana Garcia-Diaz¹

Colette Smith²

Margaret A Johnson³

Alison J Rodger^{2,4}

Sanjay Bhagani^{3,4}

Tanzina Haque¹

Daniel P. Webster^{1,3}

¹ Department of Virology, Royal Free London NHS Foundation Trust, London, UK

² Department of Infection and Population Health, University College London, London, UK

³ Department of HIV Medicine, , Royal Free London NHS Foundation Trust, London, UK

⁴ Department of Infectious Diseases, Royal Free London NHS Foundation Trust, London, UK

* Present address: Faculty of Science and Technology, University of Westminster, London, UK

"This is the author's version of a work accepted for publication by International Medical Press. Changes resulting from the publishing process, including peer review, editing and formatting, might not be reflected in this document. A definitive version was published in *Antiviral Therapy*, Vol 20, Iss 3 (2015), pp. 361-363, doi:[10.3851/imp2871](https://doi.org/10.3851/imp2871). ©2014 International Medical Press."

This final accepted version of the article has undergone peer review and contains any corrections required. The published version is available at:

<http://dx.doi.org/10.3851/imp2871>

The WestminsterResearch online digital archive at the University of Westminster aims to make the research output of the University available to a wider audience. Copyright and Moral Rights remain with the authors and/or copyright owners.

Whilst further distribution of specific materials from within this archive is forbidden, you may freely distribute the URL of WestminsterResearch: (<http://westminsterresearch.wmin.ac.uk/>).

In case of abuse or copyright appearing without permission e-mail
repository@westminster.ac.uk