

WestminsterResearch

<http://www.westminster.ac.uk/westminsterresearch>

**Determinants of Influenza and Pertussis Vaccination Uptake in
Pregnancy: A Multicenter Questionnaire Study of Pregnant
Women and Healthcare Professionals**

**Wilcox, C.R., Calvert, A., Metz, J., Kilich, E., MacLeod, R., Beadon,
K, Heath, P.T., Khalil, A., Finn, A., Snape, M.D., Vandrevala, T.,
Nadarzynski, T., Coleman, M.A. and Jones, C.E.**

This is an accepted manuscript of an article published by Wolters Kluwer in The Pediatric Infectious Disease Journal, 38 (6), pp. 625-630.

The final definitive version is available online:

<https://dx.doi.org/10.1097/INF.0000000000002242>

© 2019 Wolters Kluwer

The WestminsterResearch online digital archive at the University of Westminster aims to make the research output of the University available to a wider audience. Copyright and Moral Rights remain with the authors and/or copyright owners.

Whilst further distribution of specific materials from within this archive is forbidden, you may freely distribute the URL of WestminsterResearch: (<http://westminsterresearch.wmin.ac.uk/>).

In case of abuse or copyright appearing without permission e-mail repository@westminster.ac.uk

1 **Attitudes of pregnant women and healthcare professionals**
2 **towards clinical trials and routine implementation of**
3 **antenatal vaccination against respiratory syncytial virus: a**
4 **multi-centre questionnaire study**

5
6
7 Christopher R Wilcox MBBCh ¹, Anna Calvert MRCPh ², Jane Metz MBBS ³,
8 Eliz Kilich BM BCh ², Rachael MacLeod MBChB ³, Kirsten Beadon BSc ⁴,
9 Paul T Heath MRCPh ², Asma Khalil MD ^{5,6}, Adam Finn PhD ³, Matthew D
10 Snape MD ⁴, Tushna Vandrevala PhD ⁷, Tom Nadarzynski PhD ^{8,9}, Matthew A
11 Coleman MD ¹⁰, Christine E Jones PhD ¹¹

12
13
14 1: NIHR Clinical Research Facility, University Hospital Southampton
15 NHS Foundation Trust, Southampton, UK

16
17 2: Vaccine Institute, St George's, University of London, London, UK

18
19 3: Bristol Children's Vaccine Centre, University of Bristol, Bristol, UK

20
21 4: Oxford Vaccine Group, Department of Paediatrics, University of
22 Oxford and the NIHR Oxford Biomedical Research Centre, Oxford,
23 UK

24
25 5: Fetal Medicine Unit, Department of Obstetrics and Gynaecology,
26 St George's University Hospitals NHS Foundation Trust, London, UK

27
28 6: Vascular Biology Research Centre, Molecular and Clinical
29 Sciences Research Institute, St George's, University of London,
30 London, UK

31
32 7: School of Social and Behavioural Sciences, Faculty of Business &
33 Social Sciences, Kingston University, Kingston, London, UK

34
35 8: Department of Psychology, University of Southampton,
36 Southampton, UK

37
38 9: Department of Psychology, University of Westminster, London, UK

39
40 10: Department of Fetal Medicine, Princess Anne Hospital, University
41 Hospital Southampton NHS Foundation Trust, Southampton, UK

42
43 11: Faculty of Medicine and Institute for Life Sciences, University of
44 Southampton and University Hospital Southampton NHS Foundation
45 Trust, Southampton, UK

46
47 **Abbreviated title**

48 Attitudes to antenatal RSV vaccination: a questionnaire study

49

50 **Running title**

51 Attitudes to antenatal RSV vaccination

52

53 **Corresponding author**

- 54 - Dr Christopher Wilcox
55 - NIHR Clinical Research Facility, Southampton Centre for
56 Biomedical Research, C Level West Wing, Mailpoint 218,
57 Southampton General Hospital, Tremona Road, Southampton,
58 SO16 6DY
59 - Email: christopher.wilcox@soton.ac.uk
60 - Telephone: 02381204956
61 - Fax: 02381205023

62

63 **Keywords:** Vaccination; Pregnancy; Respiratory Syncytial Virus;
64 RSV; Clinical Trials; Attitudes

65

66 **Conflict of Interests Statement**

67 CW, AC, JM, KB, PH, AK, AF, MS and CJ are investigators for clinical
68 trials done on behalf of their respective institutions, sponsored by
69 various vaccine manufacturers, but receive no personal funding for
70 these activities.

71

73

Abstract

74

75 Introduction

76 Respiratory Syncytial Virus (RSV) is a common cause of infant
77 hospitalisation and mortality. With multiple vaccines in
78 development, we aimed to determine [1] the awareness of RSV
79 amongst pregnant women and healthcare professionals (HCPs), and
80 [2] attitudes towards clinical trials and routine implementation of
81 antenatal RSV vaccination.

82

83 Methods

84 Separate questionnaires for pregnant women and HCPs were
85 distributed within four hospitals in South England (July 2017-January
86 2018).

87

88 Results

89 Responses from 314 pregnant women and 204 HCPs (18%
90 obstetricians, 75% midwives, 7% unknown) were analyzed. Most
91 pregnant women (88%) and midwives (66%) had no/very little
92 awareness of RSV, unlike obstetricians (14%). Amongst pregnant
93 women, 29% and 75% would likely accept RSV vaccination as part
94 of a trial, or if routinely-recommended, respectively. Younger
95 women (16-24 years), those of 21-30 weeks' gestation, and with
96 experience of RSV were significantly more likely to participate in
97 trials (OR: 1.42 [1.72-9.86]; OR: 2.29 [1.22-4.31]; OR: 9.07 [1.62-
98 50.86], respectively). White-British women and those of 21-30
99 weeks' gestation were more likely to accept routinely-recommended
100 vaccination (OR: 2.16 [1.07-4.13]; OR: 2.10 [1.07-4.13]).
101 Obstetricians were more likely than midwives to support clinical
102 trials (92% vs. 68%, OR: 2.50, 1.01-6.16) and routine RSV
103 vaccination (89% vs. 79%, OR: 4.08, 1.53-9.81), as were those with
104 prior knowledge of RSV, and who deemed it serious.

105

106 **Conclusion**

107 RSV awareness is low amongst pregnant women and midwives.
108 Education will be required to support successful implementation of
109 routine antenatal vaccination. Research is needed to understand
110 reasons for vaccine hesitancy amongst pregnant women and HCPs,
111 particularly midwives.

112

114 **Introduction**

115

116 Respiratory Syncytial Virus (RSV) is the leading viral cause of lower
117 respiratory tract infection and bronchiolitis in infants, and is a major
118 cause of hospitalization and mortality worldwide¹. RSV infects more
119 than 60% of children in their first year of life, and almost 100% by
120 two years of age². The estimated case fatality ratio for children
121 hospitalized with severe RSV disease is 0.3% in industrialized
122 countries, and 2.1% in developing countries³. Severe illness often
123 occurs in children under six months⁴, particularly in those born
124 prematurely or with underlying chronic illness, and the development
125 of novel prevention and treatment strategies is an international
126 priority^{5 6}.

127

128 Antenatal vaccination is an effective means of protecting young
129 infants from infection when the period of greatest susceptibility is
130 shortly after birth⁷⁻¹⁰, and is now routinely recommended for use
131 against a number of pathogens, including tetanus, influenza and
132 pertussis¹¹. No vaccine against RSV is yet approved for routine use,
133 however a number of candidates are in development^{12 13}, one of
134 which is undergoing international phase III efficacy trials in pregnant
135 women (NCT02624947)^{11 14}. An advantage of vaccination in
136 pregnancy, rather than infancy, is that protection is afforded to
137 infants from birth and extends through the period of highest risk of
138 severe disease.

139

140 Achieving vaccine acceptance amongst pregnant women and
141 maternity healthcare professionals (HCPs) has proven to be a
142 considerable public health challenge, particularly in developed
143 countries, and uptake of routine vaccination (especially influenza)
144 remains suboptimal¹⁵. Furthermore, recruitment of pregnant
145 women into clinical trials may be difficult, particularly as historically
146 they have been excluded from participation, and there is a paucity
147 of information regarding their recruitment and retention^{16 17}. Pre-
148 emptively ascertaining the level of awareness of RSV amongst
149 pregnant women and HCPs, as well as their attitudes to vaccine
150 clinical trials and routine implementation of an RSV vaccine, may
151 allow us to identify interventions to optimise both recruitment for
152 future trials and uptake in a routine setting.

153

154 Our aims were to determine [1] the level of awareness of RSV
155 amongst pregnant women and HCPs, and [2] their attitudes towards
156 clinical trials and routine implementation of RSV antenatal
157 vaccination.

158

159

160

161

162

163

164

165

166

167

168

170 **Methods**

171

172 **Questionnaire design and development**

173 Two separate anonymized questionnaires were developed for
174 pregnant women and maternity HCPs (see supplementary
175 information). These were developed with input from a multi-
176 disciplinary study team including pediatricians, obstetricians, and
177 health psychologists. Pregnant women and maternity HCPs were
178 asked about their awareness and experience of RSV and
179 bronchiolitis, pregnant women were asked whether they would
180 hypothetically consider receiving an RSV vaccine as part of a clinical
181 trial or if a vaccine were routinely recommended, and maternity
182 HCPs if they would support clinical trials and routine
183 recommendations. Women were also asked about the number of
184 vaccines they would deem acceptable during pregnancy, and their
185 opinions regarding the design of vaccine clinical trials. Part way
186 through the questionnaire (having completed a self-assessment of
187 their prior awareness/experience of RSV and bronchiolitis),
188 participants were provided with written information on RSV and
189 bronchiolitis inside a sealed envelope. This was done in order to
190 inform further questions, whilst avoiding biasing their self-
191 assessment in the previous section. Ethical approval was granted
192 (reference 17/LO/0537) and the study was registered on
193 ClinicalTrials.gov prior to recruitment (NCT03096574).

194

195 **Study population and recruitment**

196 The questionnaire for pregnant women was administered to women
197 (aged \geq 16 years at the time of recruitment) attending for routine
198 antenatal care at four study sites in southern England: University
199 Hospital Southampton NHS Foundation Trust, Oxford University
200 Hospitals NHS Foundation Trust, University Hospitals Bristol NHS
201 Foundation Trust, and St George's University Hospitals NHS
202 Foundation Trust, London. These four study sites were selected due
203 to their high birth rates (all >4000 births/year¹⁸), and by
204 distributing our questionnaire across four hospitals we attempted to
205 increase the demographic diversity of our study population. The
206 HCP questionnaire was administered to those working in either
207 midwifery or obstetrics at the same four sites. Antenatal care for
208 low-risk women in the UK is midwife-led, with women only seeing an
209 obstetrician if they have a high-risk pregnancy, therefore the
210 majority of potential respondents to our questionnaire were
211 midwives.

212

213 Recruitment of participants took place from July 2017 to January
214 2018. Pregnant women were recruited in-person at antenatal clinics
215 and wards by members of the study team on an opportunistic (non-
216 sequential) basis over the recruitment period, and given paper
217 questionnaires to complete. For recruitment of HCPs, all
218 obstetricians and midwives at the participating institutions were
219 identified by a senior member of staff not involved in the study

220 (using email distribution lists). They were then contacted via an
221 email containing a link to an online questionnaire, followed by two
222 email reminders. Alternatively, HCPs may also have been recruited
223 in-person by the study team (in a similar fashion to pregnant
224 women), in which case they were also given paper questionnaires.
225 At the time of recruitment, information provided on the nature of
226 the questionnaire was kept to a minimum in order to avoid biasing
227 participant responses. The participant information sheet stated only
228 that the aim of the study was to better understand their attitudes
229 towards RSV and vaccination during pregnancy. Participation in the
230 study was voluntary and no financial or other incentive was offered.
231 All participants gave informed consent.

232

233 **Questionnaire data analysis**

234 Questionnaire data were entered at the lead site (Southampton) into
235 iSurvey (www.isurvey.soton.ac.uk). Statistical analysis was
236 performed using IBM SPSS Statistics version 25. Ordinal regression
237 analysis was performed, and adjusted odds ratios (ORs) and 95%
238 confidence intervals (CI) were calculated. P-values <0.05 were
239 considered as statistically significant. Multicollinearity was
240 examined using the tolerance test and the Variance Inflation Factor
241 (VIF) to ensure variables with a VIF value exceeding 2.5 were not
242 entered into the multivariate regression analysis.

243

244

247 **Results**

248

249 A total of 525 participants completed the questionnaires: 321
250 pregnant women and 204 HCPs (18% obstetricians, 75% midwives,
251 and for 7% the professional role was unknown). Seven
252 questionnaires from pregnant women, and five from HCPs, were
253 excluded due to largely incomplete or illegible responses, leaving
254 513 (98%) for analysis. The numbers of respondents were equally
255 distributed between the four study sites. The full characteristics of
256 respondents are displayed in Table 1.

257

258 **Responses from pregnant women**

259 Most pregnant women reported no (71%) or very little (17%)
260 awareness of RSV, and reported no experience (93%) [see Figure 1].
261 They were much more familiar with the term ‘bronchiolitis’ (only
262 14% had never heard of it), and bronchiolitis tended to be perceived
263 as more common and serious than RSV.

264

265 Of 312 who responded, 28% were likely/very likely, 32% not sure,
266 and 40% unlikely/very unlikely to consider receiving RSV
267 vaccination as part of a clinical trial. The most important information
268 to women was the likelihood of side effects for their baby (see
269 Figure 2). Ordinal regression analysis (see Table 2) demonstrated
270 that women were significantly more likely to accept RSV vaccination

271 as part of a clinical trial if they had direct experience of RSV (OR:
272 9.07, 95% CI: 1.62-50.86), were of younger age (16-24 years, OR:
273 1.42, 95% CI: 1.72-9.86) and of 21-30 weeks' gestation (OR: 2.29,
274 95% CI: 1.22-4.31). Women were significantly less likely to consider
275 taking part if they perceived bronchiolitis as extremely/moderately
276 serious (OR: 0.38, 95% CI: 0.15-0.93) or somewhat serious (OR:
277 0.27, 95% CI: 0.11-0.68).

278

279 More women would accept the vaccine if it was routinely
280 recommended: of 308 who responded, 40% were very likely, 35%
281 likely, 16% not sure, 5% unlikely and 4% very unlikely. Women were
282 significantly more likely to accept routine RSV vaccination if they
283 identified as White British (OR: 2.16, 95% CI: 1.22-3.83) versus non-
284 White British, and were of 21-30 weeks' gestation (OR: 2.10, 95%
285 CI: 1.07-4.13)

286

287 The most popular method of being approached regarding study
288 involvement was face-to-face by their midwife (37%), but 26%
289 wouldn't have a preference (see Figure 3). The amount of time
290 pregnant women would need to consider whether or not to
291 participate in a trial was variable, but 72% responded \leq one week
292 (17% <24 hours, 22% 1-2 days, 33% 3-7 days, 18% 2-3 weeks, and
293 10% >1 month). For the majority (82%), their decision to participate
294 wouldn't be altered if the study was a randomised controlled trial,
295 but 15% would be less likely to take part, and 3% would be more

296 likely. For 66%, their decision wouldn't be altered if the study
297 involved different doses of vaccine, but 31% would be less likely to
298 take part, and 3% would be more likely. The number of vaccines in
299 pregnancy deemed acceptable by women was variable, however
300 25% would accept two vaccines or less, 27% would accept three,
301 11% four, 6% five, and 32% would accept more than five (i.e. as
302 many as were recommended). Finally, in the free-text comments
303 (see supplementary information), some women raised concerns
304 regarding side-effects for their baby, and others stated support for
305 vaccination, often describing personal experience.

306

307 **Responses from maternity healthcare professionals**

308 HCPs had greater awareness and experience of RSV than pregnant
309 women, however obstetricians were significantly more familiar than
310 midwives with both RSV (OR: 9.42, 95% CI: 5.08-25.30, p<0.0001)
311 and bronchiolitis (OR 2.68, 95% CI: 1.29-5.55, p=0.008) [see Figure
312 1].

313

314 Of 192 HCPs who responded, 72% were likely/very likely, 19% not
315 sure, and 9% unlikely/very unlikely to support a clinical trial of RSV
316 vaccination. The most important information to HCPs was the
317 likelihood of side effects for the baby. Ordinal regression analysis
318 (see Table 2) demonstrated that HCPs were significantly more likely
319 to consider supporting a clinical trial if they were obstetricians (OR:
320 2.50, 95% CI: 1.01-6.16), had good/some understanding of RSV (OR:

321 4.42, 95% CI: 1.10-17.83), and perceived RSV as extremely (OR:
322 4.85, 95% CI: 1.11-21.28) or moderately/somewhat serious (OR:
323 4.16, 95% CI :1.26-13.75). Likelihood of support also varied between
324 study sites, with HCPs from sites A, B and C being significantly more
325 likely to support a trial than those in site D.

326

327 More HCPs would support administration of the vaccine if it was
328 routinely recommended: 47% definitely, 34% likely, 14% not sure,
329 4% unlikely and 0.5% very unlikely. Obstetricians were significantly
330 more likely than midwives to support the administration of a routine
331 RSV vaccine (OR: 4.08, 95% CI: 1.53-9.81), as were those HCPs with
332 good/some understanding of RSV (OR: 6.07, 95% CI: 1.23-29.93)
333 and those who perceived RSV as moderately/somewhat serious (OR:
334 4.41, 95% CI: 1.32-14.78) [see Table 3]. Likelihood of supporting a
335 routine RSV vaccine also varied significantly by study site with HCPs
336 from sites A, B being significantly more likely to support routine
337 vaccination than those in site D. Finally, in the free-text comments
338 [see supplementary information] some HCPs reported concerns
339 regarding the possibility of side-effects for the baby.

340

341

342

343

344

345

347 **Discussion**

348

349 The high burden of RSV infection has driven recent efforts to
350 develop an effective antenatal vaccine. This is a large multi-centre
351 study in which we have attempted to establish the level of
352 awareness of RSV, and attitudes to vaccine clinical trials and routine
353 implementation of an RSV vaccine during pregnancy.

354

355 The awareness of RSV was low amongst pregnant women and
356 midwives, compared with obstetricians. Younger pregnant women,
357 those of 21-30 weeks' gestation, and those recalling direct
358 experience of RSV, were significantly more likely to consider
359 involvement in an RSV vaccine trial; and direct face-to-face
360 interaction with a midwife was the preferred method of potential
361 recruitment (amongst those who had a preference). Encouragingly,
362 the majority of women would accept routine RSV vaccination, yet
363 some (25%) would still be unsure or unlikely to accept vaccination,
364 particularly those of ethnic minorities, and one-quarter would accept
365 ≤ 2 vaccines during pregnancy. Approximately 70% and 80% of
366 HCPs would be likely to support an RSV vaccine trial and routine
367 RSV vaccination respectively. Obstetricians were more likely than
368 midwives to support both RSV trials and routine vaccination, as
369 were those with prior knowledge of RSV and those who perceived it
370 as a serious cause of infection. Support for potential RSV trials and
371 routine vaccination also varied significantly by study site.

372

373 It is notable that the awareness of RSV is so low given that RSV-
374 associated respiratory tract infection is one of the commonest
375 causes of infant hospitalisation and mortality worldwide ¹. Being
376 thoroughly informed as to the indication and efficacy of vaccination
377 has been shown to significantly increase the probability of its
378 acceptance^{19 20}. Therefore, with a number of RSV vaccine
379 candidates currently in development, further education of both
380 pregnant women and HCPs will be needed if we are to optimise
381 engagement with vaccination trials and eventual uptake of RSV
382 vaccines as part of routine care. Both pregnant women and HCPs
383 seemed to better identify with the term bronchiolitis than RSV, and
384 therefore specifically highlighting the link between these may be
385 helpful in educational strategies. We do note that those who
386 perceived bronchiolitis as serious were significantly less likely to
387 consider participating in an RSV trial, however it is possible that this
388 is a result of confounding due to a lack of knowledge regarding
389 bronchiolitis. It is also interesting to note that women of 21-30
390 weeks' gestation were significantly more accepting of both RSV
391 trials and routine vaccination, perhaps due to a sense of
392 reassurance following their 20-week anomaly scan and subsequent
393 clinical review. Finally, the finding that women of ethnic minorities
394 were less likely to accept routine RSV vaccination has been similarly
395 observed in a number of previous studies of routinely-recommended
396 vaccines²¹⁻²³, yet the underlying reasons remain poorly understood,

397 and may include cultural/religious differences, as well as language
398 barriers.

399

400 It is concerning that a number of the HCPs surveyed in this study
401 would be unlikely to support either clinical trials or routine
402 vaccination against RSV. Maternity HCPs can be strong advocates
403 for antenatal vaccination, and encouragement from them
404 (particularly midwives) may increase intention by up to 20 times²⁴²⁵.

405 Furthermore, HCPs are well-placed to facilitate clinical trial
406 recruitment by identifying and speaking directly to eligible women,
407 and addressing specific concerns about research safety and
408 practicality¹⁷. It is important to note that obstetricians were
409 significantly more willing to provide support for both clinical trials
410 and routine vaccination than midwives, independent of their prior
411 knowledge/experience of RSV or bronchiolitis. Barriers to
412 engagement of midwives and nurses in research that have been
413 identified in previous studies, include high workload, insufficient
414 staff numbers and resources, a lack of confidence, and a lack of a
415 research-supportive culture^{26 27}. Finally, the observed differences in
416 support for both routine vaccination and clinical trials between
417 study sites also suggests that there may be a potential risk of health
418 inequalities based on differing recommendations across the South of
419 England. All four sites had been involved in trials of antenatal
420 vaccination (including RSV trials) prior to this study, and all have
421 recently embedded vaccination into their routine antenatal care

422 service. Site D only recently set up this vaccination service however
423 (following the completion of this study), whereas it has been
424 operating at the other sites for a longer period of time. They also
425 report having comparatively less involvement from clinical teams in
426 their vaccination trials. This may therefore, at least in part, explain
427 the lower acceptance at this institution compared with sites A, B and
428 C.

429

430 **Implications for clinical practice and research**

431 It is clear that education about RSV and bronchiolitis for pregnant
432 women will be required in order to optimise uptake rates of
433 antenatal RSV vaccination if it is introduced into routine care. Such
434 education should highlight the safety and benefits of vaccination for
435 their child, as studies have consistently shown that perception of
436 potential harm to the baby is the primary reason for vaccine refusal^{25 28}, whereas messages emphasising the protective benefits
437 conferred to infants is a major motivator for pregnant women to
438 undergo vaccination²⁹. As well as face-to-face counselling, possible
439 strategies could include paper and online education resources^{40 30},
440 as well as mobile phone text messages (such as Text4baby³¹) and
441 smart phone apps (such as MatImmms³²). Education for HCPs on RSV
442 and bronchiolitis will also be required in order to ensure active
443 promotion of vaccination, and individual institutions should aim to
444 tackle any general vaccine hesitancy within their own working body.

446

447 With regards to improving uptake into future antenatal vaccine
448 trials, it is important to note that the majority of our respondents
449 wouldn't be deterred by a randomized controlled trial design, and
450 that direct face-to-face interaction with an HCP was the preferred
451 method of recruitment. Improving study team outreach and forming
452 integrated networks between research teams and healthcare
453 providers/clinical staff may help improve clinicians' willingness to
454 promote clinical studies to their patients, as well as pregnant
455 women's willingness to join studies ¹⁷, and this has proven a
456 successful method of recruiting pregnant women in previous studies
457 ^{33 34}. Social media and web-based recruitment may be used as a
458 cost-effective supplement to traditional recruitment methods, and
459 facilitate participation of traditionally harder-to-reach populations ¹⁷
460 ³⁵, however this approach may be less successful for higher-risk
461 intervention-based studies, including antenatal vaccine trials.

462

463 Finally, it should be noted that there are other potential antenatal
464 vaccines in development (including group B streptococcus and
465 cytomegalovirus ¹¹), for which education and support from staff will
466 also be required for successful implementation ²² . Furthermore, it is
467 also worth considering that whilst a third of our respondents would
468 accept as many vaccines as were recommended, many women may
469 be reluctant to accept high numbers of vaccines, especially if given
470 on separate occasions^{36 37}. Pragmatic research is therefore required

471 to consider the logistical aspects of future antenatal vaccine
472 delivery.

473

474 **Strengths and limitations**

475 This study had significant numbers of respondents, and by
476 distributing our questionnaire across four hospitals in southern
477 England we attempted to maximise the diversity of our study
478 population. That said, the responses to the questionnaire cannot be
479 taken as representative of all pregnant women and maternity HCPs.
480 Our respondents were all recruited from antenatal clinics based in
481 tertiary hospitals, and therefore it is also possible that our sample
482 was missing subsets of the population that tend to be more anti-
483 vaccination. Future studies might benefit from recruiting over a
484 wider geographical area, and from different types of sites (such as
485 non-tertiary hospitals and primary care), and perhaps utilising
486 online recruitment via pregnancy-associated websites and social
487 media. It may have been also beneficial to collect socio-economic
488 data from our participants in order to assess the representativeness
489 of our study sample. Other limitations are that data on the uptake of
490 antenatal vaccination was not collected from women's medical
491 records following delivery, and data on the uptake of influenza
492 vaccination amongst HCPs wasn't collected. Finally, the number of
493 pregnant women/HCPs approached, and the number who declined
494 participation (as well as their reasons for doing so) was not
495 recorded, and we are therefore unable to report this data.

496

497 **Conclusions**

498 RSV awareness appears low amongst pregnant women and
499 midwives in the UK. Education will be required to optimise
500 engagement with vaccination trials and eventual uptake of RSV
501 vaccination following routine implementation, with an emphasis on
502 women of ethnic minorities. Active promotion of vaccination must
503 be incorporated into routine antenatal care, and further research is
504 needed to understand reasons for vaccine hesitancy amongst both
505 pregnant women and HCPs, particularly midwives.

506

507

508

510 **Figure captions [images to be reproduced in colour**
511 **online only]:**

512
513 **Figure 1:** Reported familiarity and experience with RSV (A & B) and
514 bronchiolitis (C) amongst pregnant women, midwives and
515 obstetricians, prior to their involvement in this study.

516
517 **Figure 2:** Information that would be considered most important to
518 the pregnant women in this study when deciding whether to take
519 part in a research study of an RSV vaccine (A), and other factors
520 which would discourage them from taking part (B).

521
522 **Figure 3:** Preferred method of being approached regarding
523 potential clinical trial involvement amongst the pregnant women in
524 this study

525
526 **Acknowledgements**

527
528 The authors would also like to thank all the pregnant women and
529 healthcare staff who took part in the questionnaire, Stephen Yekini
530 for his assistance with data collection in Southampton, and all of the
531 non-study staff that helped facilitate recruitment in the participating
532 sites.

533
534 **Author Contributions**

535 CW drafted the manuscript and was principal investigator. All
536 authors contributed to questionnaire design and critically revised
537 the manuscript. CW, AC, JM, EK, RM, KB, PH, AK, AF, MS, TV, TN, MC
538 and CJ were involved in study set up and data collection at the
539 participating sites. CW, TN and CJ performed the data analysis. CJ
540 conceived the study and was chief investigator. All authors
541 approved the final version of the manuscript.

542
543 **Conflict of Interests Statement**

544 CW, AC, JM, KB, PH, AK, AF, MS and CJ are investigators for clinical
545 trials done on behalf of their respective institutions, sponsored by
546 various vaccine manufacturers, but receive no personal funding for
547 these activities.

548
549 **Funding**

550 The study was supported by a grant from the British Paediatric
551 Allergy Immunity and Infection Group (BPAIIG). BPAIIG had no role in
552 the study design, data collection, data analysis/interpretation,
553 report writing, or the decision to submit the manuscript for
554 publication.

555
556 **Clinical trial registration**

557 The questionnaire study was registered on ClinicalTrials.gov prior to
558 recruitment (NCT03096574).

559

560 **Ethical approval**

561 Ethical approval was granted from the West London & GTAC NHS
562 Research Ethics Committee (reference 17/LO/0537).

564

565

- ## References
- 566 1. Hall CB, Weinberg GA, Iwnae MK, Blumkin AK, Edwards KM, et
567 al. The Burden of Respiratory Syncytial Virus Infection in Young
568 Children. *New Engl J Med* **360**, 588-598 (2009).
 - 569 2. Resch, B. Burden of respiratory syncytial virus infection in
570 young children. *World J Clin Pediatr* **1**, 8-12 (2012).
 - 571 3. Nair H, Nokes DJ, Gessner BD, Dherani M, Madhi SA, Singleton
572 RJ. Global burden of acute lower respiratory infections due to
573 respiratory syncytial virus in young children: a systematic review
574 and meta-analysis. *Lancet* **375**, 1545-55 (2010).
 - 575 4. Nokes DJ, Ngama M, Bett A, Abwao J, Munywoki P, English M,
576 et al. Incidence and Severity of Respiratory Syncytial Virus
577 Pneumonia in Rural Kenyan Children Identified through Hospital
578 Surveillance. *Clin Infect Dis* **49**, 1341-1349 (2009).
 - 579 5. Yamin D, Jones FK, DeVincenzo JP, Gertler S, Kobiler,
580 Townsend JP, et al. Vaccination strategies against respiratory
581 syncytial virus. *Proc Natl Acad Sci USA*. **113**, 13239-13244 (2016).
 - 582 6. Saso, A. & Kampmann, B. Vaccination against respiratory
583 syncytial virus in pregnancy: a suitable tool to combat global infant
584 morbidity and mortality? *Lancet Infect Dis* **16**, e153-e163 (2016).
 - 585 7. Poehling KA, Szilagyi PG, Staat MA, et al. Impact of maternal
586 immunization on influenza hospitalizations in infants. *Am J Obstet
587 Gynecol.* 2011;204(6 Suppl 1):S141-8.
 - 588 8. Amirthalingam G, Andrews N, Campbell H, et al. Effectiveness
589 of maternal pertussis vaccination in England: an observational
590 study. *Lancet.* 2014;384(9953):1521-1528.
 - 591 9. Zaman K, Roy E, Arifeen SE, et al. Effectiveness of maternal
592 influenza immunization in mothers and infants. *New Engl J Med.*
593 2008;359(15):1555-64.
 - 594 10. Madhi SA, Cutland CL, Kuwanda L, et al. Influenza Vaccination
595 of Pregnant Women and Protection of Their Infants. *New Engl J Med.*
596 2014;371(10):918-931.
 - 597 11. Jones CE, Calvert A, Le Doare K. Vaccination in Pregnancy –
598 Recent Developments. *The Pediatric Infectious Disease Journal*
599 (2018). doi:10.1097/INF.0000000000001822
 - 600 12. Munoz FM, Piedra PA, Glezen WP. Safety and immunogenicity
601 of respiratory syncytial virus purified fusion protein-2 vaccine in
602 pregnant women. *Vaccine* **21**, 3465-7 (2003).
 - 603 13. Chu HY, Steinhoff MC, Magaret A, Zaman K, Roy E, Langdon G,
604 et al. Respiratory Syncytial Virus Transplacental Antibody Transfer
605 and Kinetics in Mother-Infant Pairs in Bangladesh. *J Infect Dis* **210**,
606 1582-1589 (2014).
 - 607 14. Kaaijk P, Luytjes W, Rots N. Vaccination against RSV: is
608 maternal vaccination a good alternative to other approaches? *Hum
609 Vaccin immunother* **9**, 1263-7 (2013).
 - 610 15. Bisset K, Paterson P. Strategies for increasing uptake of
611 vaccination in pregnancy in high-income countries: A systematic
612 review. *Vaccine* (2018). doi:10.1016/j.vaccine.2018.04.013

- 613 16. Blehar MC, Spong C, Grady C, Goldkind SF, Sahin L, Clayton JA.
614 Enrolling pregnant women: issues in clinical research. *Womens
615 Health Issues* **23**, e39–45 (2013).
- 616 17. Frew PM, Saint-Victor DS, Brewinski Isaacs M, Kim S, Swamy
617 GK, Sheffield JS. Recruitment and retention of pregnant women into
618 clinical research trials: an overview of challenges, facilitators, and
619 best practices. *Clin Infect Dis.* **59 Suppl 7**, S400–7 (2014).
- 620 18. NHS Digital. NHS Maternity Statistics, England 2016-2017.
621 (2018).
- 622 19. Donaldson B, Jain P, Holder BS, Lindsey B, Regan L,
623 Kampmann B. What determines uptake of pertussis vaccine in
624 pregnancy? A cross sectional survey in an ethnically diverse
625 population of pregnant women in London. *Vaccine* **33**, 5822–5828
626 (2015).
- 627 20. Panda B, Stiller R, Panda, A. Influenza vaccination during
628 pregnancy and factors for lacking compliance with current CDC
629 guidelines. *J Matern Fetal Neonatal Med* **24**, 402–406 (2011).
- 630 21. Goldfarb IT, Little S, Brown J, Riley LE. Use of the combined
631 tetanus-diphtheria and pertussis vaccine during pregnancy. *AJ
632 Obstet Gynecol* **211**, 299.e1–299.e5 (2014).
- 633 22. McQuaid F, Jones C, Steven Z, Meddaugh G, O'Sullivan C,
634 Donaldson B, et al. Antenatal vaccination against Group B
635 streptococcus: Attitudes of pregnant women and healthcare
636 professionals in the UK to participation in clinical trials and routine
637 implementation. *Acta Obstet Gynecol Scand* **97**, 330–340(2018).
- 638 23. Dempsey A, Brewer S, Sevick C, et al. Tdap vaccine attitudes
639 and utilization among pregnant women from a high-risk population.
640 *Hum Vaccin Immunother* **12**, 872–878 (2016).
- 641 24. Wiley KE, Massey PD, Cooper SC, Wood NJ, Ho J, et al. Uptake
642 of influenza vaccine by pregnant women: a cross-sectional survey.
643 *Med J Aust* **198**, 373–375 (2013).
- 644 25. Wiley KE, Massey PD, Cooper SC, Wood NJ, Quinn HE, Leask J.
645 Pregnant women's intention to take up a post-partum pertussis
646 vaccine, and their willingness to take up the vaccine while pregnant:
647 A cross sectional survey. *Vaccine* **31**, 3972–3978 (2013).
- 648 26. Mitchell K, Baillie L, Phillips N. Increasing nurse and midwife
649 engagement in research activity. *Nursing Standard* **29**, 37–42
650 (2015).
- 651 27. Purkis J, Jackson JA, Hundt G, Stockman C. Increasing nursing
652 research capacity in the workplace. *Nurs Times* **104**, 28–31 (2008).
- 653 28. Sim J, Ulanika A, Katikireddi S, Gorman D. 'Out of two bad
654 choices, I took the slightly better one': Vaccination dilemmas for
655 Scottish and Polish migrant women during the H1N1 influenza
656 pandemic. *Public Health* **125**, 505–511 (2011).
- 657 29. Marsh H, Malik F, Shapiro E, Omer S, Frew P. Message Framing
658 Strategies to Increase Influenza Immunization Uptake Among
659 Pregnant African American Women. *Matern Child Health J* **18**, 1639–
660 1647 (2014).
- 661 30. Zhou L, Zhang D, Yang C, Wang Y. Harnessing social media for

- 662 health information management. *Electron Commer R A* **27**, (2018).
- 663 31. Gazmararian JA, Elon L, Yang B, Graham M, Parker R.
- 664 Text4baby program: an opportunity to reach underserved pregnant
- 665 and postpartum women? *Matern Child Health J* **18**, 223–32 (2014).
- 666 32. Holder B, Borqeau M, Donaldson B, Davies J, Kampmann B.
- 667 MatImmms: A smartphone app to inform and educate women about
- 668 maternal immunisation. Poster presentation at 4th International
- 669 Neonatal and Maternal Immunisation Symposium, Brussels,
- 670 Belgium, 10-12/09/2017
- 671 33. Coleman-Phox K, Laraia BA, Adler N, Vieten C, Thomas M, Epel
- 672 E. Recruitment and retention of pregnant women for a behavioral
- 673 intervention: lessons from the maternal adiposity, metabolism, and
- 674 stress (MAMAS) study. *Prev Chronic Dis* **10**, (2013).
- 675 34. Kerver JM, Elliot MR, Norman GS, Sokol RJ, Keating DP,
- 676 Copeland GE, et al. Pregnancy recruitment for population research:
- 677 the National Children's Study vanguard experience in Wayne
- 678 County, Michigan. *Paediatr Perinat Epidemiol* **27**, 303–11 (2013).
- 679 35. Shere M, Zhao AY, Koren G. The role of social media in
- 680 recruiting for clinical trials in pregnancy. *PLoS ONE* **9**, e92744
- 681 (2014).
- 682 36. Webb H, Street J, Marshall H. Incorporating immunizations into
- 683 routine obstetric care to facilitate Health Care Practitioners in
- 684 implementing maternal immunization recommendations. *Hum*
- 685 *Vaccin Immunother* **10**, 1114–21 (2014).
- 686 37. Krishnaswamy S, Wallace E, Buttery J, Giles M. Strategies to
- 687 implement maternal vaccination: A comparison between standing
- 688 orders for midwife delivery, a hospital based maternal immunisation
- 689 service and primary care. *Vaccine* **36**, 1796–1800 (2018).
- 690
- 691
- 692
- 693
- 694

Table 1: Characteristics of the questionnaire respondents (pregnant women and maternity healthcare professionals)

Characteristic	Pregnant women, n=314	Healthcare professionals, n=199
Age		
16-24	34 (11%)	
25-30	107 (34%)	
31-35	92 (29%)	
36-40	58 (19%)	
41+	13 (4%)	
Gestation (weeks)		
<12	8 (2%)	
12-16	37 (12%)	
17-20	31 (10%)	
21-30	55 (18%)	
31-36	93 (30%)	
>37	76 (24%)	
Study site		
A	88 (28%)	43 (22%)
B	77 (25%)	53 (27%)
C	79 (25%)	61 (31%)
D	70 (22%)	42 (21%)
Ethnicity		
Asian (British, Indian, Pakistani, Bangladeshi, Chinese, other)	25 (8%)	4 (2%)
Black (British, African, Caribbean, other)	17 (5%)	4 (2%)
White (British, Irish, other)	248 (79%)	175 (88%)
Mixed (Caribbean, African, Asian, other)	11 (4%)	6 (3%)
Other ethnic group (Arab, other)	3 (1%)	0 (0%)
Did not want to say	1 (0.3%)	1 (1%)
No response	10 (3%)	9 (5%)
Has children		
No	142 (45%)	72 (36%)
Yes	172 (55%)	127 (64%)
Profession		
Obstetrics		37 (19%)
Midwifery		151 (76%)
No response		11 (6%)
Midwifery seniority		
Band 5 (newly-qualified midwife)	8 (5%)	
Band 6 (junior midwife)	84 (56%)	
Band 7 (senior midwife)	46 (30%)	
Band 8 (midwifery manager)	8 (5%)	
No response	5 (3%)	
Obstetrician seniority		
Specialty training years 1-3 (or equivalent)	8 (22%)	
Specialty training years 4-6 (or equivalent)	6 (16%)	
Specialty training years 7-8 (or equivalent)	6 (16%)	
Consultant	17 (46%)	
Time spent working in maternity care (years)		
<2	17 (9%)	
2-5	29 (15%)	
6-10	37 (19%)	
11-15	20 (10%)	
16-20	26 (13%)	
>21	62 (31%)	
No response	8 (4%)	

703 **Table 2:** Ordinal regression analysis of factors predicting pregnant women's willingness to
 704 consider undergoing RSV vaccination during pregnancy as part of a clinical trial, or if routinely
 705 recommended

Variable	Number who'd be 'extremely likely' or 'likely' to accept RSV vaccination as part of a clinical trial	Adjusted odds ratio (95% CI)	Number who'd be extremely likely' or 'likely' to accept RSV vaccination if routinely recommended	Adjusted odds ratio (95% CI)
Age in years				
16-24	18/34 (53%)	1.42 (1.72-9.86) **	27/34 (79%)	0.68 (0.28-1.67)
25-35	54/199 (27%)	1.18 (0.67-2.07)	149/198 (75%)	0.71 (0.39-1.28)
36-45	16/70 (23%)	1.00 for reference	53/70 (76%)	1.00 for reference
Gestation in weeks				
<12	3/8 (38%)	1.99 (0.46-8.51)	6/8 (75%)	0.67 (0.15-3.00)
12-20	24/68 (35%)	1.26 (0.72-2.22)	52/68 (76%)	1.17 (0.65-2.10)
21-30	18/55 (33%)	2.29 (1.22-4.31) **	42/54 (78%)	2.10 (1.07-4.13) *
31+	43/168 (26%)	1.00 for reference	128/168 (76%)	1.00 for reference
Study site				
Site A	25/86 (29%)	0.80 (0.40-1.59)	65/87 (75%)	0.99 (0.49-2.00)
Site B	23/77 (30%)	0.72 (0.35-1.49)	62/76 (82%)	1.26 (0.59-2.69)
Site C	20/79 (25%)	0.54 (0.26-1.10)	55/76 (72%)	0.78 (0.37-1.63)
Site D	20/70 (29%)	1.00 for reference	50/69 (72%)	1.00 for reference
Previous children				
Yes	50/171 (29%)	1.13 (0.71-1.81)	122/171 (71%)	0.64 (0.39-1.05)
No	39/141 (28%)	1.00 for reference	110/137 (80%)	1.00 for reference
Ethnicity				
White British	66/224 (29%)	1.27 (0.73-2.21)	177/223 (79%)	2.16 (1.22-3.83) **
Non-White British	23/88 (26%)	1.00 for reference	55/85 (65%)	1.00 for reference
Previous RSV experience				
Direct experience	5/8 (63%)	9.07 (1.62-50.86) *	8/8 (100%)	8.20 (0.71-94.16)
Indirect experience	5/13 (38%)	1.11 (0.32-3.81)	10/13 (77%)	1.09 (0.30-3.96)
No experience	79/291 (27%)	1.00 for reference	214/287 (75%)	1.00 for reference
RSV familiarity				
Good/some understanding	5/14 (36%)	0.54 (0.12-2.30)	11/14 (79%)	1.77 (0.37-8.56)
Poor understanding	21/77 (27%)	0.80 (0.47-1.38)	55/76 (72%)	0.96 (0.55-1.68)
No understanding	63/219 (29%)	1.00 for reference	164/216 (76%)	1.00 for reference
Perceived RSV frequency				
Extremely/moderately common	18/50 (36%)	1.12 (0.53-2.35)	39/51 (76%)	1.03 (0.47-2.23)
Somewhat common	34/99 (34%)	1.52 (0.88-2.61)	75/98 (77%)	0.93 (0.53-1.64)
Slightly/not at all common	37/143 (26%)	1.00 for reference	107/141 (76%)	1.00 for reference
Perceived RSV severity				
Extremely/moderately serious	43/129 (33%)	1.22 (0.58-2.57)	100/129 (78%)	1.31 (0.60-2.86)
Somewhat serious	33/117 (28%)	0.93 (0.47-1.84)	87/115 (76%)	1.06 (0.52-2.18)
Slightly/not at all serious	13/43 (30%)	1.00 for reference	33/43 (77%)	1.00 for reference
Bronchiolitis familiarity and experience				
Good/moderate understanding and direct/indirect experience	27/88 (31%)	1.30 (0.65-2.60)	68/89 (76%)	0.75 (0.36-1.53)
Slight understanding	29/102 (28%)	1.13 (0.63-2.00)	77/101 (76%)	0.81 (0.44-1.48)
No understanding	32/120 (27%)	1.00 for reference	86/116 (74%)	1.00 for reference
Perceived bronchiolitis frequency				
Extremely/moderately common	33/107 (31%)	0.67 (0.33-1.37)	85/107 (79%)	1.04 (0.49-2.19)
Somewhat common	26/96 (27%)	1.25 (0.68-2.31)	69/95 (73%)	1.36 (0.72-2.60)
Slightly/not at all common	26/101 (26%)	1.00 for reference	73/98 (74%)	1.00 for reference
Perceived bronchiolitis severity				
Extremely/moderately serious	55/190 (29%)	0.38 (0.15-0.93) *	143/188 (76%)	0.63 (0.24-1.65)
Somewhat serious	19/84 (23%)	0.27 (0.11-0.68) *	62/84 (74%)	0.52 (0.20-1.36)
Slightly/not at all serious	11/28 (39%)	1.00 for reference	20/26 (77%)	1.00 for reference

706 *= $p<0.05$; **= $p<0.01$

709 **Table 3:** Ordinal regression analysis of factors predicting the willingness of healthcare
 710 professionals to support RSV vaccination during pregnancy as part of a clinical trial, or if
 711 routinely recommended

Variable	Number who'd be 'very likely' or 'likely' to support RSV vaccination as part of a clinical trial	Adjusted odds ratio (95% CI)	Number who'd be 'very likely' or 'likely' to support RSV vaccination if routinely recommended	Adjusted odds ratio (95% CI)
Professional group				
Obstetrics	34/37 (92%)	2.50 (1.01-6.16) *	33/37 (89%)	4.08 (1.53-9.81) **
Midwifery	102/151 (68%)	1.00 for reference	119/151 (79%)	1.00 for reference
Time in maternity care				
21+ years	46/62 (74%)	0.51 (0.14-1.83)	46/62 (74%)	0.43 (0.12-1.62)
11-20 years	31/46 (67%)	0.38 (0.11-1.34)	34/46 (74%)	0.79 (0.22-2.86)
2-10 years	47/66 (71%)	0.68 (0.22-2.10)	60/66 (91%)	1.39 (0.43-4.42)
<2 years	14/17 (82%)	1.00 for reference	15/17 (88%)	1.00 for reference
Study site				
Site A	30/41 (73%)	3.94 (1.46-10.61) **	34/41 (83%)	3.95 (1.39-11.26) *
Site B	35/53 (66%)	3.19 (1.23-8.30) *	46/53 (87%)	6.23 (2.22-17.46) ***
Site C	51/61 (84%)	5.80 (2.36-14.21) ***	47/61 (77%)	1.97 (0.81-4.83)
Site D	22/37 (59%)	1.00 for reference	29/37 (78%)	1.00 for reference
Has own children				
Yes	88/127 (69%)	0.59 (0.28-1.24)	101/127 (80%)	0.86 (0.39-1.91)
No	50/65 (77%)	1.00 for reference	55/65 (85%)	1.00 for reference
Ethnicity				
White British	126/175 (72%)	1.01 (0.34-3.06)	142/175 (81%)	1.41 (0.44-4.46)
Non-White British	12/17 (71%)	1.00 for reference	14/17 (82%)	1.00 for reference
RSV experience				
Direct experience	22/26 (85%)	2.65 (0.79-8.86)	24/26 (92%)	1.41 (0.39-5.07)
Indirect experience	20/27 (74%)	1.17 (0.42-3.31)	23/27 (85%)	0.74 (0.25-2.22)
No experience	96/139 (69%)	1.00 for reference	109/139 (78%)	1.00 for reference
RSV familiarity				
Good/some understanding	19/22 (86%)	4.42 (1.10-17.83) *	20/22 (91%)	6.07 (1.23-29.93) *
Poor understanding	87/114 (76%)	1.81 (0.88-3.73)	91/114 (80%)	1.07 (0.51-2.24)
No understanding	32/55 (58%)	1.00 for reference	44/55 (80%)	1.00 for reference
Perceived RSV frequency				
Extremely common	29/36 (81%)	1.43 (0.45-4.51)	30/36 (83%)	1.96 (0.57-6.76)
Moderately/somewhat common	84/116 (72%)	0.92 (0.43-1.98)	95/116 (82%)	1.20 (0.54-2.67)
Slightly/not at all common	25/39 (64%)	1.00 for reference	30/39 (77%)	1.00 for reference
Perceived RSV severity				
Extremely serious	27/35 (77%)	4.85 (1.11-21.28) *	26/35 (74%)	1.25 (0.28-5.55)
Moderately/somewhat serious	113/138 (82%)	4.16 (1.26-13.75) *	117/138 (85%)	4.41 (1.32-14.78) *
Slightly/not at all serious	8/17 (47%)	1.00 for reference	12/17 (71%)	1.00 for reference
Bronchiolitis familiarity and experience				
Good/moderate understanding and indirect/direct experience	58/77 (75%)	0.84 (0.10-6.94)	66/77 (86%)	0.99 (0.12-8.35)
Slight understanding	78/111 (70%)	0.98 (0.13-7.49)	87/111 (78%)	0.98 (0.13-7.56)
No understanding	2/4 (50%)	1.00 for reference	3/4 (75%)	1.00 for reference
Perceived bronchiolitis frequency				
Extremely common	29/34 (85%)	1.05 (0.34-3.25)	27/34 (79%)	0.55 (0.17-1.80)
Moderately/somewhat common	88/124 (71%)	1.44 (0.63-3.29)	104/124 (84%)	1.07 (0.45-2.51)
Slightly/not at all common	21/34 (62%)	1.00 for reference	25/34 (74%)	1.00 for

Perceived bronchiolitis severity				reference
<i>Extremely serious</i>	36/47 (77%)	0.35 (0.054-2.28)	38/47 (81%)	0.96 (0.15-6.39)
<i>Moderately/somewhat serious</i>	96/136 (71%)	0.29 (0.052-1.65)	111/136 (82%)	0.54 (0.10-2.99)
<i>Slightly/not at all serious</i>	6/9 (67%)	1.00 for reference	7/9 (78%)	1.00 for reference

712

713

714 *=p<0.05; **=p<0.01; ***=p<0.001

716

Supplementary information

717

718

719

1) Questions for pregnant women analysed in this study

720

(1) Before taking part in this survey, how familiar were you with Respiratory Syncytial Virus (sometimes shortened to RSV)?

- I have never heard of it
- I have heard of it, but don't really know what it is
- I know some facts about what it is
- I have a good understanding about RSV infection and its implications

727

(2) What experience do you have of RSV?

- I have no experience of it
- I know someone who has experience of it
- I have direct experience of it

732

(3) How common do you think RSV infection is in babies and young children?

- Not at all common
- Slightly common
- Somewhat common
- Moderately common
- Extremely common

739

(4) How serious do you think RSV infection is for babies and young children?

- Not at all serious
- Slightly serious
- Somewhat serious
- Moderately serious
- Extremely serious

746

(5) Before taking part in this survey, how familiar were you with bronchiolitis in babies and young children?

- I have never heard of it
- I have heard of it but don't know what it is
- I know some facts about it
- I know what it is and know someone who has experience of it
- I know what it is and have direct experience of it

754

(6) How common do you think bronchiolitis is in babies and young children?

- Not at all common
- Slightly common
- Somewhat common
- Moderately common
- Extremely common

761

762

(7) How serious do you think bronchiolitis is for babies and young children?

- Not at all serious
- Slightly serious
- Somewhat serious
- Moderately serious
- Extremely serious

769

770

(8) Would you be *potentially* willing to receive a RSV vaccine during pregnancy as part of a research study to determine its safety and effectiveness, before the vaccine is approved for routine use?

773
774 Your response to this question will not affect whether or not you receive further information about such
775 studies and **does not mean** that you are agreeing to take part in any vaccine research studies.
776

- 777 Extremely unlikely
778 Unlikely
779 Neutral/not sure
780 Likely
781 Extremely likely
782

783 **(9) What information would you consider to be important when considering taking part in a**
784 **research study of a RSV vaccine?**

785
786 **Please rank the top 3 most important to you: (1= most important information for you to know)**

- 787
788 How common RSV is
789 How serious RSV is
790 Number of healthy adults who have received the vaccine
791 Number of pregnant women who have received the vaccine
792 Likelihood of side effects for me
793 Likelihood of side effects for my baby
794

795 **(10) One type of a research study is a “Randomised Controlled Trial” where there are two (or**
796 **more) groups who are treated exactly the same, except only one group gets the true vaccine**
797 **under investigation. The other group may get a ‘placebo’ (dummy or inactive) injection.**

798
799 This type of study allows the researchers to check that any differences between the groups are
800 due to the vaccine only. Importantly, patients or staff do not get to choose whether they receive
801 the proper vaccine or the dummy.

802
803 **After reading the above information:**

- 804 I would be less likely to take part as I would want to guarantee that I would have the vaccine
805 I would be more likely to take part as I might not get the vaccine
806 This would not affect my decision
807

808
809 **(11) In some randomised controlled trials, patients are given different doses (amounts) of the**
810 **vaccine under investigation in order to work out which is the best dose to use in future vaccines.**
811 **These different doses would be calculated before the trial starts, but patients or staff involved in**
812 **the study do not get to choose which of these doses they receive.**

813 **After reading the above information:**

- 814 I would be less likely to take part
815 I would be more likely to take part
816 This would not affect my decision
817

818 **(12) What other factors would discourage you from taking part in a research study of a vaccine**
819 **in pregnancy?**

820 **Please rank the following: (1=factor that would most discourage you, 4=factor least likely to**
821 **discourage you)**

- 822 Number of hospital visits
823 Number of home visits
824 Number of blood tests for me
825 Number of blood tests for baby
826 Other, please
827 specify.....
828 ...
829

830 **(13) How would you most like to be approached about taking part in a research study?**

831 **Tick one answer:**

- 832 Asked by my midwife

- 833 Asked by my obstetrician
 834 Asked by my GP
 835 Given a leaflet/poster with contact details for the study team
 836 Adverts of the internet (e.g. pregnancy forums)
 837 Email from the study team
 838 Approached directly by the study midwife/doctor
 839 I wouldn't mind how I was approached
 840
 841 Other:.....
 842
 843
(14) If you were approached about taking part in a research study, how much time would you like to fully consider whether or not you would like to take part?
 844 <24 hours
 845 1-2 days
 846 3-7 days
 847 2-3 weeks
 848 >1 month
 849
(15) Would you be willing to receive this vaccine in pregnancy if it was routinely recommended for use in pregnancy in the NHS?
 850 Definitely
 851 Probably
 852 Maybe
 853 Probably not
 854 Definitely not
 855
(16) There are a number of different vaccines that are being designed for use in pregnancy to protect mothers and infants against severe infection. How many vaccines would be acceptable to you in pregnancy?
 856 0
 857 1
 858 2
 859 3
 860 4
 861 5
 862 More than 5
 863
(27) How old are you in years?
 864 16-24 25-30 31-35 36-40 41-45 46+
 865
(28) How many weeks pregnant are you?
 866 Less than 12 12-16 17-20 21-30 31-36 37+
 867
(19) To what ethnic group do you feel you belong? (Please circle)
 868

White British - English / Welsh / Scottish / Northern Irish / British Irish - Gypsy or Irish Traveller specify)..... - Other (please specify)	Black / African / Caribbean / Black - African - Caribbean - Other (please
---	---

 869

Mixed/Multiple ethnic groups - White and Black Caribbean - White and Black African specify)..... - White and Asian - Other (please specify)	Other ethnic group - Arab - Other (please
---	--

 870

- 952 Somewhat common
953 Moderately common
954 Extremely common
955

956 **(7) How serious do you think bronchiolitis is for young children?**

- 957 Not at all serious
958 Slightly serious
959 Somewhat serious
960 Moderately serious
961 Extremely serious
962

963 **(8) Would you be potentially willing to support a randomised controlled trial of RSV vaccine in**
964 **pregnancy to determine its safety and how well it prevents infection in children, by signposting**
965 **the study to women?**

966 *Your response to this question will not affect whether or not you receive further information about such*
967 *studies*

- 968
969 Extremely unlikely
970 Unlikely
971 Neutral/not sure
972 Likely
973 Extremely likely
974

975 **(9) Would you be willing to support the administration of this vaccine if it was routinely**
976 **recommended for use in the NHS?**

- 977 Definitely
978 Probably
979 Maybe
980 Probably not
981 Definitely not
982

983 **(10) What factors would influence your decision regarding whether or not you would be willing**
984 **to support involvement in a RSV vaccine research study before it is licensed?**

- 985
986 **Please rank the top 3 factors: (1=factor that would most influence you)**
987
988 The number of pregnant women who had previously received the vaccine in research studies
989 How common RSV is in children
990 Seriousness of RSV infection in young children
991 How effective the vaccine is in preventing *RSV infection*
992 How effective the vaccine is in preventing *severe RSV disease*
993 Risk of side effects for the mother
994 Risk of side effects for developing baby
995 Other (please specify):
996

997
998 **(11) How many pregnant women would the vaccine have to be safely tested on in a research**
999 **study for you to consider supporting such a trial?**

- 1000 None
1001 Over 10
1002 Over 100
1003 Over 500
1004 Over 1000
1005 Over 5000
1006 Over 10,000
1007 I would not support such a trial
1008

1009 **(12) Which healthcare professional group do you belong to?**

- 1010 Obstetrics
1011 Midwifery

1012	<input type="checkbox"/> Other (please state)	
1013	
1014		
1015	(13) How long have you worked in maternity care?	
1016	<input type="checkbox"/> Under 2 years	
1017	<input type="checkbox"/> 2-5 years	
1018	<input type="checkbox"/> 6-10 years	
1019	<input type="checkbox"/> 11-15 years	
1020	<input type="checkbox"/> 16-20 years	
1021	<input type="checkbox"/> 21+ years	
1022		
1023	(14) What is your grade?	
1024	<i>1. Midwifery/nursing staff</i>	
1025	Band 4 <input type="checkbox"/> Band 5 <input type="checkbox"/> Band 6 <input type="checkbox"/> Band 7 <input type="checkbox"/> Band 8 <input type="checkbox"/> Band 9 <input type="checkbox"/>	
1026	<i>2. Obstetricians</i>	
1027	ST 1-3 (or equivalent) <input type="checkbox"/> ST 4-6 (or equivalent) <input type="checkbox"/> ST 7-8 (or equivalent) <input type="checkbox"/>	
1028	<input type="checkbox"/>	
1029		
1030	(15) Have you had any children before?	
1031	<input type="checkbox"/> Yes.	
1032	If yes, how many?.....	
1033	What are their ages?	
1034	Child 1: Less than 1 <input type="checkbox"/> 1-5 <input type="checkbox"/> 6-10 <input type="checkbox"/> 11-16 <input type="checkbox"/> 17+ <input type="checkbox"/>	
1035	Child 2: Less than 1 <input type="checkbox"/> 1-5 <input type="checkbox"/> 6-10 <input type="checkbox"/> 11-16 <input type="checkbox"/> 17+ <input type="checkbox"/>	
1036	Child 3: Less than 1 <input type="checkbox"/> 1-5 <input type="checkbox"/> 6-10 <input type="checkbox"/> 11-16 <input type="checkbox"/> 17+ <input type="checkbox"/>	
1037	Child 4: Less than 1 <input type="checkbox"/> 1-5 <input type="checkbox"/> 6-10 <input type="checkbox"/> 11-16 <input type="checkbox"/> 17+ <input type="checkbox"/>	
1038	<input type="checkbox"/> No	
1039		
1040	(16) To what ethnic group do you feel you belong? (Please circle)	
1041		
1042	White	Black / African / Caribbean / Black
1043	British	
1044	- English / Welsh / Scottish / Northern Irish	- African
1045	/ British Irish	- Caribbean
1046	- Gypsy or Irish Traveller	- Other (please
1047	specify).....	
1048	- Other (please specify)	
1049		
1050	Mixed/Multiple ethnic groups	Other ethnic group
1051	- White and Black Caribbean	- Arab
1052	- White and Black African	- Other (please
1053	specify).....	
1054	- White and Asian	
1055	- Other (please specify)	
1056		
1057	Asian / Asian British	I'd prefer not to say
1058	- Indian	
1059	- Pakistani	
1060	- Bangladeshi	
1061	- Chinese	
1062	- Other (please specify)	
1063		
1064	(17) Optional: Do you have any comments or concerns about vaccination or vaccine research	
1065	studies during pregnancy?	
1066		
1067		
1068		
1069		

1070
1071
1072
1073
1074 3) **Free-text comments from pregnant women and healthcare**
1075 **professionals**

1076
1077 *Response to the question: Do you have any **comments** or **concerns** about vaccination or vaccine*
1078 *research studies during pregnancy?*

1079
1080 **Pregnant women**

- 1081 1. I think vaccine trials are very risky even though very important so every available
1082 information should be made available to the participant before commencing including
1083 all known possible side effects
1084
1085 2. Many vaccines contain unsafe levels of mercury in some cases some are produced
1086 on human tissue (DNA) and contain various other toxins. I believe a baby is born with
1087 a perfect immune system which takes up to 3 years to fully develop and that it's not
1088 healthy injecting a perfectly healthy child with chemicals and toxins (mercury)
1089
1090 3. I am glad to hear that the NICE guidelines will be reviewed and that possibly new
1091 vaccines will be introduced
1092
1093 4. I am taking part in a RSV vaccine trial
1094
1095 5. I'm very keen for my baby to have as many vaccines as possible & fully support such
1096 research
1097
1098 6. I would want the vaccine fully tested and approved before I would have it
1099
1100 7. Our daughter suffered from bronchiolitis at age 2 weeks old so as long as the vaccine
1101 was safe we would definitely have it to prevent this baby suffering like our daughter
1102 did
1103
1104 8. I would consider vaccination if I was having a normal singleton pregnancy
1105
1106 9. I'm a bit of a unique case because I've had an adverse reaction to a vaccine in the
1107 past and wouldn't risk it in pregnancy unless I had to
1108
1109 10. Child died at 20 months. RSV sounds very like what my son had when he died
1110
1111 11. No concerns. I am very pro vaccinations both for myself during pregnancy and for my
1112 children
1113
1114 12. I am having a slightly bumpy pregnancy and this is one of the reasons I would be
1115 reluctant to take part in a research study which could increase the risks for the
1116 pregnancy complications. If I was a low-risk person I would be more willing to take
1117 part. Likewise, if this wasn't my first baby I might be more willing
1118
1119 13. Information about the potential side effects of the trial vaccinations would have been
1120 helpful for me to make more informed decisions
1121
1122 14. I've not heard of RSV before sounds concerning and something I would have liked to
1123 have been told about earlier in my pregnancy
1124
1125 15. I've heard of many children developing chest infections as young babies and anything
1126 to avoid this I feel should be actively encouraged
1127

- 1128
1129 16. I would like the opportunity to ask more questions and have more information before
1130 agreeing to vaccination
1131
1132 17. I would only have medication in pregnancy that has been approved by the BMA.
1133 Diabetics have a lot of complications anyway
1134
1135 18. No - thank you for all the amazing work/research you do
1136
1137 19. I believe the stage of drug trial to be more pertinent to the decision-making process
1138 than the number of vaccinations received.
1139
1140 20. My concern in taking part in a research study is the unknown side effects to my baby
1141 and whether the potential side effects would cause more harm than the virus itself.
1142 Whilst I appreciate research needs to be done and the vaccine will have been
1143 thoroughly tested on other test groups testing pregnant women/babies is still a
1144 concern for me
1145
1146 21. Not really aware enough of the issue to comment on some of the questions
1147
1148 22. In my experience, the flu vaccine has made me ill. I would not feel comfortable having
1149 a trial vaccine as a first-time mother

1150
1151 **Maternity healthcare professionals**

- 1152
1153 1. I would want to see safety data in non-pregnant participants concerning side effects
1154 and efficacy before I supported vaccine studies on pregnant women. I understand
1155 that effectiveness in preventing baby bronchiolitis could not be assessed using non-
1156 pregnant subjects but would reassure health workers that we aren't supporting an
1157 action that could cause harm.
1158
1159 2. I would worry about safety /side effects to mum and baby if not tested before being
1160 given to pregnant women
1161
1162 3. Knowledge to midwives about RSV is very limited without having first-hand
1163 experience of it or working alongside paediatric teams. It's not widely taught in
1164 training perhaps because our care for infants doesn't go much beyond 10-28 days
1165 postnatally
1166
1167 4. My son needed ECMO because of this infection but he was too unstable to transfer to
1168 Gt O S we very nearly lost him at 12 days old. He caught it from his sister who was 2
1169 and poorly when he was born. This serious infection wiped the first 2 months of a
1170 normal newborn period for us. He did get asthma as a child and took months to catch
1171 up.
1172
1173 5. Side effects - baby especially.
1174
1175 6. I'm not convinced that RS virus needs vaccination. Depends on the severity of
1176 chance of later disease in the child. I think we build up immunity ourselves and
1177 therefore the number needed to treat is probably high to prevent severe RS virus
1178 infection in children.
1179
1180 7. I would want some evidence that the vaccine is safe.
1181
1182 8. My children were born at 27/40 and 32/40 week's gestation. For our 27/40 week-old
1183 baby it was very serious.
1184
1185 9. Potential risks to unborn and ability to be honest with mother about risks v benefits.
1186
1187 10. Risk to unborn.

- 1188
1189 11. Can't really answer question of how many women vaccines would have to be safely
1190 tested on as I don't know what the predicted rate of adverse reactions/side effects. As
1191 long as sufficiently powered I would be happy. No concerns as long as properly
1192 conducted. Vaccination research environment is heavily regulated so very confident.
1193
- 1194 12. That the vaccine is safe for the mother and unborn child. This has to be paramount
1195 and is of high concern with the majority of the public.
1196
- 1197 13. When testing for side effects -there should be follow-up of at least 5 years on the
1198 child whose mother received the vaccine. We are woefully short on long-term effects
1199 and in order to fully discuss (and understand) the effects of vaccinations in pregnancy
1200 these time-frames should be mandatory. Lack of long-term data does not reassure
1201 me that we should be vaccinating in pregnancy.
1202
- 1203 14. Effect on the baby that are so far unknown. Another vaccine could it be combined
1204 with present vaccines?
1205
- 1206 15. I would worry about a trial re the long term unknown effects on the health of children
1207 whose mothers received the vaccine whilst they were in utero.
1208
- 1209 16. Yes, the potential risks to mother and unborn baby
1210