

WestminsterResearch

<http://www.westminster.ac.uk/westminsterresearch>

Environmental heterogeneity of Staphylococcus species from alkaline fermented foods and associated toxins and antimicrobial resistance genetic elements

Ouoba, L.I.I., Voudibio-Mbozo, A.B., Anyogu, A., Obioha, P.I., Lingani-Sawadogo, H., Sutherland, J.P., Jespersen, L. and Ghoddusi, H.B.

NOTICE: this is the authors' version of a work that was accepted for publication in International Journal of Food Microbiology. Changes resulting from the publishing process, such as peer review, editing, corrections, structural formatting, and other quality control mechanisms may not be reflected in this document. Changes may have been made to this work since it was submitted for publication. A definitive version was subsequently published in International Journal of Food Microbiology, DOI: 10.1016/j.ijfoodmicro.2019.108356, 2019.

The final definitive version in International Journal of Food Microbiology is available online at:

<https://dx.doi.org/10.1016/j.ijfoodmicro.2019.108356>

© 2019. This manuscript version is made available under the CC-BY-NC-ND 4.0 license <https://creativecommons.org/licenses/by-nc-nd/4.0/>

The WestminsterResearch online digital archive at the University of Westminster aims to make the research output of the University available to a wider audience. Copyright and Moral Rights remain with the authors and/or copyright owners.

Whilst further distribution of specific materials from within this archive is forbidden, you may freely distribute the URL of WestminsterResearch: (<http://westminsterresearch.wmin.ac.uk/>).

In case of abuse or copyright appearing without permission e-mail repository@westminster.ac.uk

1 **Environmental heterogeneity of *Staphylococcus* species from alkaline**
2 **fermented foods and associated toxins and antimicrobial resistance genetic**
3 **elements**

4
5 L. Irène I. Ouoba^{a,b*}, Alain Brice Voudibio-Mbozo^c, Amarachukwu Anyogu^d,
6 Promiselynda I. Obioha^a, Hagrétou Lingani-Sawadogo^e, Jane P. Sutherland^a, Lene
7 Jespersen^f, and Hamid B. Ghoddusi^a

8 ^a Microbiology Research Unit, School of Human Sciences, London Metropolitan
9 University, UK

10 ^b Ouoba-Consulting, London, UK

11 ^c Faculté des Sciences, Université Marien N’Gouabi, Brazzaville, Congo

12 ^d Department of Life Sciences, Faculty of Science and Technology, University of
13 Westminster, UK

14 ^e Département Technologie Alimentaire/IRSAT/CNRST, Ouagadougou, Burkina Faso

15 ^f Microbial Ecology and Food Fermentation, Faculty of Sciences, University of
16 Copenhagen

17 **Corresponding author:** Dr L. Irène I. Ouoba, Microbiology Research Unit, School of
18 Human Sciences, London Metropolitan University London N7 8DB, UK

19 Tel.: +44 (0) 20 71334154, Email: ouobairene@hotmail.com

20 **Abstract**

21 Different samples of three products including Bikalga and Soumbala from
22 Burkina Faso (West Africa) and Ntoba Mbodi from Congo-Brazzaville (Central Africa)
23 were evaluated. The bacteria (400) were phenotyped and genotypically characterized
24 by Rep-PCR, PFGE, 16S rRNA and *rpoB* gene sequencing and *spa* typing. Their
25 PFGE profiles were compared with those of 12,000 isolates in the Center for Disease
26 Control (CDC, USA) database. They were screened for the production of enterotoxins,
27 susceptibility to 19 antimicrobials, presence of 12 staphylococcal toxin and 38 AMR
28 genes and the ability to transfer erythromycin and tetracycline resistance genes to
29 *Enterococcus faecalis* JH2-2.

30 Fifteen coagulase negative (CoNS) and positive (CoPS) species characterised
31 by 25 Rep-PCR/PFGE clusters were identified: *Staphylococcus arlettae*, *S. aureus*, *S.*
32 *cohnii*, *S. epidermidis*, *S. gallinarum*, *S. haemolyticus*, *S. hominis*, *S. pasteurii*, *S.*
33 *condimenti*, *S. piscifermentans*, *S. saprophyticus*, *S. sciuri*, *S. simulans*, *S. warneri*
34 and *Macrococcus caseolyticus*. Five species were specific to Soumbala, four to
35 Bikalga and four to Ntoba Mbodi. Two clusters of *S. gallinarum* and three of *S. sciuri*
36 were particular to Burkina Faso. The *S. aureus* isolates exhibited a *spa* type t355 and
37 their PFGE profiles did not match any in the CDC database. Bacteria from the same
38 cluster displayed similar AMR and toxin phenotypes and genotypes, whereas clusters
39 peculiar to a product or a location generated distinct profiles. The toxin genes
40 screened were not detected and the bacteria did not produce the staphylococcal
41 enterotoxins A, B, C and D. AMR genes including *blazA*, *cat501*, *dfr(A)*, *dfr(G)*, *mecA*,
42 *mecA1*, *msr(A)* and *tet(K)* were identified in CoNS and CoPS. Conjugation
43 experiments produced JH2-2 isolates that acquired resistance to erythromycin and
44 tetracycline, but no gene transfer was revealed by PCR.

45 The investigation of the heterogeneity of *Staphylococcus* species from alkaline
46 fermented foods, their relationship with clinical and environmental isolates and their
47 safety in relation to antimicrobial resistance (AMR) and toxin production is anticipated
48 to contribute to determining the importance of staphylococci in alkaline fermented
49 foods, especially in relation to the safety of the consumers.

50

51

52 **Key words:** Alkaline fermented foods; *Staphylococcus* spp.; diversity; distribution;
53 safety; toxins; AMR

54

Accepted Manuscript

55 **1. Introduction**

56 Alkaline fermented foods are products that are generally rich in proteins (30-
57 50%) and their production involves a fermentation process associated with an increase
58 of pH to values as high as 10.0 (Parkouda et al., 2009; Voudibio Mbozo et al., 2017).
59 They are mainly known in Africa and Asia, where they constitute low cost sources of
60 protein foods and play an important part in the diet of people (Parkouda et al., 2009).
61 It is generally recognized that the pH rise is the result of accrued degradation of raw
62 material proteins that leads to the release of peptides, essential amino acids as well
63 as alkalinizing substances such as ammonia (Kiers et al., 2000). In some cases, alkali-
64 treatments during the production are responsible for the pH augmentation (Parkouda
65 et al., 2008).

66 Bikalga, Soumbala, and Ntoba Mbodi are alkaline fermented foods which are
67 produced through the traditional fermentation of roselle seeds (*Hibiscus sabdariffa*),
68 African locust bean seeds (*Parkia biglobosa*) and cassava leaves (*Manihot esculenta*
69 Crantz) respectively. Production of Bikalga and Soumbala comprises several
70 processing stages of the seeds: cleaning, boiling (about 95°C for 24-40 h), dehulling
71 (Soumbala), fermentation (48-72 h), steaming (Bikalga) and drying (Ouoba et al.,
72 2008a; Parkouda et al., 2008; Parkouda et al., 2009). They are the most popular
73 traditional fermented foods used as condiments in Burkina Faso (West Africa) and are
74 also produced in other African countries. Ntoba Mbodi, which is popular in the Republic
75 of the Congo (Central Africa), is made of cassava leaves and the production process
76 does not involve heating steps. After being cleaned and cut, the leaves are distributed
77 into small portions in larger leaves, wrapped and left to ferment for 2-4 days (Voudibio
78 Mbozo et al., 2017). The fermented cassava leaves are then used to prepare sauces,
79 which are eaten with starch-based foods. In Bikalga, Soumbala and Ntoba Mbodi, pH

80 values up to 8.5, 9 and 10 have been respectively reported (Ouoba, 2015; Ouoba,
81 2017; Parkouda et al., 2009; Voudibio Mbozo et al., 2017).

82 The key microorganisms involved in alkaline fermentations are *Bacillus* species
83 such as *B. subtilis*, *B. licheniformis*, *B. amyloliquefaciens*, *B. pumulis*, and *B.*
84 *xylanilyticus* that can degrade proteins, lipids, and carbohydrates of the raw materials
85 and bring about the nutritional quality and safety of the product (Parkouda et al., 2009).
86 However, secondary microorganisms such as lactic acid bacteria, *Staphylococcus*
87 spp. and various Gram-positive and Gram-negative bacteria are often isolated with no
88 clear indication of their role in the fermentation. The production techniques of most
89 alkaline fermented foods are traditional, leading to variability of microbial population,
90 nutritional quality, product stability and potential safety issues. It is believed that some
91 of the secondary microorganisms are in fact contaminants from the producers,
92 processing materials and environments that have poor hygienic quality in most
93 traditional fermentation settings. Some secondary microorganisms are part of the
94 original microbiota of the raw materials that are suppressed by products released
95 during the fermentation. For example, those that are alkali sensitive will be eliminated
96 or reduced during the fermentation.

97 *Staphylococci* are Gram-positive non-motile cocci ubiquitously spread in
98 nature, and are commonly isolated from humans, animals, food and environmental
99 sources. More than 70 species and subspecies of the *Staphylococcus* genus have
100 been characterized so far (LPSN, 2018) with *Staphylococcus aureus* being one of the
101 most studied species due to its clinical importance. *Staphylococcus* species are
102 usually divided into two groups i.e. coagulase-positive (CoPS) and coagulase-
103 negative (CoNS) according to their ability to coagulate rabbit plasma. Various
104 *Staphylococcus* species have been recurrently isolated in many alkaline and non-

105 alkaline vegetable based fermented foods, but their role in such fermentation is
106 unknown. However, their importance in the fermentation of meat-based products has
107 been well documented. They occur in the production of various types of sausages
108 where they constitute a part of the autochthonous microbiota or are added as starter
109 cultures in combination with lactic acid bacteria (Fonseca et al., 2013; Milicevic et al.,
110 2014). *Staphylococcus equorum*, *S. saprophyticus*, *S. xylosus* and *S. carnosus* are
111 the most common species found in fermented sausages, but other species such as *S.*
112 *warneri*, *S. vitulinus*, *S. pasteurii*, *S. epidermidis*, *S. lentus*, *S. haemolyticus*, *S.*
113 *intermedius*, and *S. hominis* can also be occasionally isolated (Milicevic et al., 2014).
114 They have been reported to be involved in biochemical activities during meat
115 fermentation such as the degradation of proteins, lipids and peroxide as well as the
116 reduction of nitrite (Casaburi et al., 2008; Milicevic et al., 2014). These activities
117 generate compounds such as peptides, amino acids, carbonyls and volatile
118 compounds that contribute to the nutritional quality, specific organoleptic properties
119 and safety of the fermented meat products (Milicevic et al., 2014).

120 Although beneficial properties of *Staphylococcus* species in food have been
121 documented, many species are most known worldwide for causing food poisoning and
122 other types of clinical infections in humans and animals including e.g. bacteraemia,
123 wound infections, pyogenic lesions, and mastitis. Moreover, they present detrimental
124 antimicrobial resistance (AMR) profiles that make some of their infections hard to treat.
125 With regard to toxin production in food, staphylococcal food poisoning is one of the
126 most common food-borne diseases. It results from the ingestion of heat stable
127 staphylococcal enterotoxins (SEs) produced in food by enterotoxigenic strains of
128 mainly *S. aureus* (Hennekinne et al., 2012). Symptoms related to ingestion of such
129 toxins are fast-acting, occurring generally within 30 min to 8 h and lasting for about 24

130 hours. Individuals generally experience vomiting, nausea, stomach cramps, and
131 diarrhoea. Severe illnesses related to staphylococcal food poisoning are rare but
132 occasionally, it can be deadly, especially in children, elderly and patients weakened
133 by a long-term illness. *Staphylococcus aureus* is considered one of the world's leading
134 causes of disease outbreaks associated with food consumption. Other CoPS such as
135 *S. intermedius* and CoNS such as *S. epidermidis*, *S. sciuri*, *S. simulans*, *S. xylosum*,
136 *S. equorum*, *S. lentus* and *S. capitis* have been reported to harbour genes coding for
137 SEs and to produce some SEs (Casaes Nunes et al., 2015; Crass and Bergdoll, 1986).
138 However, information on their real implication in staphylococcal food poisoning outbreaks
139 is scarce (Breckinridge and Bergdoll, 1971; Podkowik et al., 2013). Significantly, the
140 incidence of AMR has continually increased in staphylococci, as well as in other types
141 of bacteria. This is attributed to factors such as the frequent antibiotic administration
142 to livestock, or misuse and overuse of antimicrobials. Therefore, the food chain
143 constitutes one possible route of AMR transfer (Angulo et al., 2004). Both CoPS and
144 CoNS from food origin exhibit a wide range of AMR genetic elements coding for
145 resistance to different antimicrobials such as methicillin, tetracycline, penicillin, and
146 vancomycin (Gundogan et al., 2005; Simeoni et al., 2008). The most frequently
147 encountered resistant staphylococcus is methicillin-resistant *Staphylococcus aureus*
148 (MRSA), which is resistant to multiple antimicrobials and is responsible for hard-to-
149 treat infections. Thus, it constitutes a serious public health concern, as it causes
150 thousands of deaths per year worldwide and constitutes a global economic burden.

151 The study aimed to screen the environmental heterogeneity of *Staphylococcus*
152 species from alkaline fermented foods, their relationship with clinical isolates
153 worldwide and their safety with regards to genes encoding antimicrobial resistance

154 (AMR) and toxin production. This is anticipated to elucidate their distribution and
155 importance in such foods, especially in relation to the safety of the consumers.

156 **2. Materials and Methods**

157 2.1 Identification of the bacteria

158 2.1.1 Phenotypic characterisation

159 The bacteria (400) were isolated from different samples of Bikalga (9), Ntoba Mbodi
160 (8) and Soumbala (10) collected at different markets and production sites in Burkina
161 Faso (Bikalga and Soumbala) and the Republic of the Congo (Ntoba Mbodi),
162 from areas constituting major consumers of these food products. Each sample (10 g)
163 was homogenised for 1 min in 90 ml of maximum recovery diluent (MRD, Oxoid) and
164 ten-fold dilutions (10^{-2} - 10^{10}) prepared. The dilutions (0.1 ml) were spread on nutrient
165 agar (NA; Oxoid CM0003, Basingstoke, UK) for the enumeration of total aerobic
166 mesophilic bacteria. The plates were incubated aerobically at 37°C and the count
167 recorded after 2 days. For enumeration and isolation of the presumptive
168 *Staphylococcus* population, the dilutions were spread on Baird Parker (BPA; Oxoid)
169 and Mannitol Salt (MSA; Oxoid) agars and the plates incubated aerobically at 37°C for
170 2 days. Bacteria isolated from the two the latter media were purified by repeated
171 streaking on NA, maintained in nutrient broth (NB; Oxoid CM001) containing 20% (v/v)
172 glycerol (Sigma G8773, Gillingham, UK) as well as on cryobeads and stored at -20°C
173 for further analyses.

174 For the phenotypic characterization, the isolates were streaked on NA and
175 examined for colony and cell morphology, as well as tested for Gram, catalase,
176 oxidase and coagulase reactions. The Gram reaction was evaluated using the KOH
177 method (Gregersen, 1978) and the oxidase test using *Pseudomas fragi* as positive
178 control and *Brochothrix thermosphacta* as negative control. The coagulase test was

179 performed using the Staphylase test kit (Oxoid DR0595) and following the
180 manufacturer's instructions. The cell morphology was determined by light microscopy
181 (Nikon Model Eclipse, E400, Japan). The phenotypic screening resulted in the
182 elimination from the study of 310 non-staphylococcal isolates which were able to grow
183 on BPA and MSA and whose colony morphologies initially suggested they were
184 potential *Staphylococcus* spp. Thus, 90 isolates were retained for further
185 characterization.

186 2.1.2 Molecular typing

187 2.1.2.1 DNA extraction

188 The bacteria were grown on Brain Heart Infusion Agar (BHI-A) prepared by
189 mixing BHI (Oxoid CM1135) and technical agar (Oxoid LP0013, 15g/L). After
190 incubation at 37°C for 48 h, the DNA was extracted from purified cultures following the
191 manufacturer's instructions using InstaGene Matrix (Bio-Rad 732-6030, Hemel
192 Hempstead, UK).

193 2.1.2.2 Repetitive element palindromic-polymerase chain reaction (rep-PCR)

194 The isolates were characterized by rep-PCR using the GTG5 primer as
195 described by Ouoba et al. (2008a). Bacteria showing the same DNA profile were
196 clustered in the same group by visual screening and cluster analysis using the
197 Bionumerics system (2.50, Dice's Coefficient of similarity, Unweighted pair group
198 method with arithmetic: UPGMA; Applied Maths, Saint-Martens-Latem, Belgium).

199 2.1.2.3 Pulse Field Gel Electrophoresis (PFGE)

200 In order to assess the worldwide distribution of the types of *S. aureus* found in
201 the fermented foods, PFGE of SmaI-digested genomic DNA was performed according
202 to the method proposed by the Center for Disease Control and Prevention/Pulse Net
203 (CDC, USA) as described by Van Balen et al. (2013). The Salmonella serotype

204 Branderup strain H9812 digested with XbaI was used as a molecular size marker. The
205 PFGE profiles of the *Staphylococcus* isolates were compared with those in a CDC
206 database that contains worldwide profiles of about 12,000 *S. aureus* strains recovered
207 from outbreaks, surveillance, reference and the environment, and submitted to the
208 database.

209 2.1.2.4 Sequencing of the 16S rRNA and rpoB genes

210 The bacteria were identified by the amplification and sequencing of the 16S
211 rRNA and *rpoB* genes. For the 16S rRNA gene amplification, the method and the pA
212 and pE primers described by Ouoba et al. (2008a) were used. For the amplification
213 of the *rpoB* gene, primers 2491F and 3554R described by Drancourt and Raoult (2002)
214 were used with the following PCR mixture: 5 µl of 10 X PCR buffer containing 15 mM
215 of MgCl₂ (Applied Biosystems N8080160), 0.5 µl of dNTP (1.25 mM), 0.5 µl of each
216 primer (21 pmol/µl), 0.2 µl of AmpliTaq polymerase (5 U; Applied Biosystems N808-
217 0160) and 41.3 µl of sterile high purity water. The PCR conditions were as follows:
218 initial denaturation at 94°C for 2 min, 35 cycles of 94°C for 30 s, 52°C for 30 s, 72°C
219 for 1 min and a final extension step at 72°C for 5 min.

220 After the amplification of both genes, positive amplicons were checked by
221 electrophoresis and purified using the QIAquick PCR purification kit (Qiagen, Crawley,
222 UK). The 16S rRNA and *rpoB* genes were sequenced (SourceBioscience, Cambridge,
223 UK) using primers pD (3.2 pmol/µl) and 2491F/ 3554R (3.2 pmol/µl) respectively. The
224 16S rRNA gene sequences were analysed in the ezbiocloud server (Yoon et al., 2017)
225 whereas those of the *rpoB* gene were screened in the GenBank/EMBL/DDBJ
226 database (National Center for Biotechnology, MD, USA).

227 2.1.2.4 *Spa* typing

228 A multiplex PCR method and primers recommended by the European
229 Reference Laboratory for antimicrobial resistance (EURL-AR, 2012), and that allows
230 at the same time the detection of the *spa* gene and the *S. aureus* methicillin resistance
231 genes *mecA*, *mecC* was used. The method includes the use of a PCR mixture
232 containing 2.5 µl of 2xGreen PCR Master Mix (Thermo Scientific K1081), 2 µl (10 µM)
233 of a mixture of the forward primers, 2 µl (10 µM) of a mixture of the reverse primers
234 and 6.5 µl of high purity water. The PCR conditions described by Stegger et al. (2012)
235 were applied.

236 2.2 Screening haemolysis and toxin production potential of the bacteria

237 2.2.1 Haemolytic activity on blood agar

238 The *Staphylococcus* isolates were screened for their haemolytic activity on
239 blood agar as follows: Columbia agar base (Oxoid CM003) was autoclaved at 121 °C
240 for 15 min and sheep blood (5%; Oxoid SR0051B) added after cooling to 50 °C, before
241 distribution into Petri dishes. The isolates were then streaked on the agar and
242 incubated at 37 °C for 48 h. Haemolysis was identified by the appearance of a zone
243 of clearing around the colonies.

244 2.2.2 Production of staphylococcal enterotoxins A, B, C and D

245 The ability of the bacteria to produce the staphylococcal enterotoxin A, B, C
246 and D was investigated using the staphylococcal enterotoxin reversed passive latex
247 agglutination (SET-RPLA) toxin detection kit (Oxoid TD0900). Each isolate was grown
248 at 37°C for 24h in 10 ml of Tryptone Soya broth (TSB; Oxoid CM0129) and 1ml of the
249 culture centrifuged using a benchtop centrifuge at 3000 rpm for 20 min. The presence
250 of toxins was screened in the supernatant according to the manufacturer's instructions.

251 2.2.3 Detection of staphylococcal enterotoxin, exfoliative and toxic shock syndrome 252 toxin genes

253 The isolates were tested for the presence of the staphylococcal enterotoxin
254 genes *sea*, *seb*, *sec*, *sed*, *see*, *seg*, *seh*, *sei*, *sej*, exfoliative toxin genes *eta* and *etb*,
255 and toxic shock syndrome toxin gene *tsst-1*. Primers described by Johnson et al.
256 (1991), Omoe et al. (2002) and Nashev et al. (2004) were used. The PCR mixture was
257 that used by Ouoba et al. (2008b). The PCR conditions were as follows: 94°C for 2
258 min followed by 35 cycles of denaturation at 94°C for 1 min, annealing at the
259 corresponding temperature for 1 min and elongation at 72°C for 1 min. The PCR ended
260 with a final extension at 72°C for 5 min.

261 2.3 Screening the AMR phenotypic and genotypic profiles of the bacteria

262 2.3.1 Inoculum preparation

263 The isolates were grown on BHI-A and pure colonies re-suspended in 1 ml of
264 Maximum Recovery Diluent (MRD; Oxoid CM0733). A quantity of the latter suspension
265 was transferred to 10 ml of MRD to obtain a turbidity of 0.5 (10^7 - 10^8 CFU/ml) on the
266 McFarland scale using a nephelometer (TREK Diagnostic systems, West Sussex,
267 UK).

268 2.3.2 Determination of the susceptibility of the bacteria to antimicrobials

269 The Minimal Inhibitory Concentration (MIC) for 19 antimicrobials (Table 1) was
270 determined using a 96 well sensititre™ plate (EUST, TREK Diagnostic Systems Ltd,
271 East Grinstead, UK) containing varying amounts of each antimicrobial. The method
272 recommended by EURL-AR (DTU, Denmark) was applied (EURL-AR, 2009). Briefly,
273 50 µl of the inoculum was mixed with 10 ml Mueller Hinton broth (MHB; Oxoid
274 CM0405) and 50 µl of this mixture were dispensed in each well. Incubation was carried
275 out aerobically at 37°C for 24 h and visible growth recorded. The susceptibility of the
276 isolates to each antimicrobial was established using the cut-off values recorded in
277 Table 1.

278 2.3.3 Detection of AMR genes

279 The presence of 38 genes encoding resistance to chloramphenicol,
280 erythromycin, gentamycin, methicillin, penicillin, quinupristin/dalfopristin, tetracycline,
281 trimethoprim, streptomycin and vancomycin were investigated (Table 1).

282 For the detection of the genes, the PCR mixtures described by Naghizadeh
283 Raeisi et al. (2018) were used. However, for the detection of the *S. aureus* methicillin
284 resistance genes *mecA* and *mecC*, the multiplex PCR (described in 2.1.2.4) and
285 primers recommended by EURL-AR (2012) were used. For the *mecA1* gene pertaining
286 to *S. sciuri*, the same mixture (except the primers) as that used for the detection of
287 erythromycin (Naghizadeh Raeisi et al., 2018) was applied. The references of the
288 primers are depicted in Table 1. Positive control isolates (EURL-AR) for most genes
289 screened were included. Moreover, all positive amplicons were sequenced to confirm
290 the identity of the genes.

291 All PCR amplifications were performed in a thermocycler (GeneAmp PCR 2700
292 system) using the following program: initial denaturation at 94°C for 3 min, followed by
293 25 - 35 cycles of 94°C for 1 min, 45-68°C for 1 min depending on the annealing
294 temperature for the individual set of primers, 72 °C for 1 min and a final extension at
295 72 °C for 10 min. After purification, the presumptive positive amplicons were
296 sequenced using the amplification primers at a concentration of 3.2 pmol/μl and the
297 similarity of the sequence tested against published sequences in GenBank.

298 2.3.4 Screening the transferability of AMR

299 The ability of *S. warneri* 105 to transfer the *msr(A)* gene to *Enterococcus*
300 *faecalis* JH2-2 and *S. saprophyticus* A17M105 to transfer the *tet(K)* gene to the same
301 recipient was investigated by conjugation experiments according to Ouoba *et al.*
302 (2008c) and Naghizadeh Raeisi et al. (2018). The donors are resistant to erythromycin

303 (*S. warneri*) and tetracycline (*S. saprophyticus*), but sensitive to rifampicin and fusidic
304 acid. The recipient *E. faecalis* JH2-2 is resistant to rifampicin and fusidic acid and
305 sensitive erythromycin and tetracycline.

306 Inocula were prepared as described above and 10 ml of the donor suspension
307 and 1 ml of that of the recipient mixed and filtered through a sterile membrane filter
308 (0.45 µm) (Whatman Laboratory Division, Maidstone, UK) using a vacuum pump
309 (Welch Thomas 2522C-02, Skokie, Illinois, USA). The filters were incubated
310 aerobically on BHI-A at 37 °C for 48h. The growth was washed off the filter with 3 ml
311 of MRD, diluted (up to 10⁻⁵) and 100 µl of each dilution spread on agar plates
312 containing different antimicrobials. For the *msr(A)* gene, the inoculation was made on
313 BHI-RFE [rifampicin (25µg/ml), fusidic acid (25 µg/ml) and erythromycin (16 µg/ml)],
314 BHI-RF [rifampicin (25 µg/ml) and fusidic acid (25 µg/ ml)], BHI-E [Erythromycin (16
315 µg/ml)] and BHI agar plates. For the tet(K) gene transfer, the mating mixtures were
316 spread on BHI-RFT [rifampicin (25µg/ml), fusidic acid (25 µg/ml) and tetracycline (10
317 µg/ml)], BHI-RF [rifampicin (25 µg/ml) and fusidic acid (25 µg/ ml)], BHI-T [(tetracycline
318 (10 µg/ml)] and BHI agar plates. The plates were incubated at 37 °C for 7-21 days.
319 Cells of the donors and recipient that had not undergone a conjugation procedure were
320 all spread on the same type of media as those used to inoculate the mating mixtures.

321 Only recipients that had acquired resistance to erythromycin and tetracycline
322 are able to grow on BHI-RFE and BHI-RFT agar plates, respectively. Potential *E.*
323 *faecalis* JH2-2 transconjugant (PET) DNA fingerprints were compared with those of
324 the recipients and donors by rep-PCR using the GTG5 primer (Ouoba et al., 2008a).
325 In addition, their MICs for erythromycin and tetracycline were determined.

326 2.3.5 Determination of the genetic background of acquired erythromycin and
327 tetracycline resistance in the PETs

328 2.3.5.1 Extraction of DNA

329 Total DNA was extracted as described above. Plasmid DNA was extracted as
330 follows: the donors, recipient and PETs were cultured for 48 h on BHI-A. A pure colony
331 was then transferred into 10 ml BHI and incubated in a shaking water bath at 37°C for
332 12 h. Plasmid DNA was extracted using the QIAGEN Plasmid miniprep Kit (QIAGEN
333 GmbH, Hilden, Germany) according to the manufacturer's instructions.

334 2.3.5.2 Amplification of the AMR genes in the PETs

335 PCR using total and plasmid DNA were performed to detect the presence of
336 the *msr(A)* and *tet(K)* genes in the PETs. Since the *S. warneri* isolate contained also
337 the *cat501*, *blazA* and *dfr(A)* genes, these were also screened in the PETs. The use
338 of plasmid DNA aimed to determine the location of genes in the bacteria and to
339 increase the chance to detect a potential transfer in the PETs.

340 3. Results

341 3.1 Identification of the bacteria

342 The pH values of the products were 5-6 in the unfermented samples and 8-9 in
343 the fermented products. The bacterial numbers recovered (CFU/g) for each product
344 were as follows: Soumbala: 10^6 - 10^{13} CFU/g on NA, 10^3 - 10^{10} on MSA, 10^1 - 10^8 on BP;
345 Bikalga: 10^8 - 10^{11} on NA, 10^7 - 10^8 on MSA, 10^4 - 10^8 on BP; Ntoba Mbodi: 10^9 - 10^{11} on
346 NA, 10^8 - 10^{10} on MSA, 10^5 - 10^8 on BP. Out of 27 samples analysed, *Staphylococcus*
347 species were recovered from 18 samples. The products where *Staphylococcus*
348 isolates were not detected included mainly samples from the production steps of
349 Bikalga and Soumbala that had undergone heat treatment, but also one sample of
350 each of the three foods obtained from different markets. As mentioned earlier, many
351 bacteria recovered on MSA and BP did not possess typical staphylococcal
352 characteristics and were not included in the advanced screening. Most of those

353 bacteria were Gram-positive (many of which were spore-forming) and Gram-negative
354 rods that were able to grow on the selective media (BPA and MSA). The 90 isolates
355 that were further screened showed variable colony and cell morphology. All were
356 cocci, arranged differently (single, pairs, clusters) according to the isolate, catalase
357 and Gram positive and oxidase negative. Two isolates from Bikalga were coagulase
358 positive and the remaining 88 bacteria were coagulase negative.

359 The microbial profile was variable according to the fermented product, the
360 production site and country (Table 2, Table 3). Genotyping by rep-PCR and PFGE
361 both generated 25 clusters representing 15 species (Table 2, Fig 1). The comparison
362 of the PFGE profiles of the *S. aureus* isolates did not generate a similarity with any of
363 the 12,000 profiles in the CDC database. By sequencing the 16S rRNA and *rpoB*
364 genes, the bacteria of eight rep-PCR/PFGE clusters were identified as *S. sciuri* (45.6
365 % of the total isolates), three as *S. gallinarum* (7.8 %), two as *S. cohnii* (4.4%), two as
366 *S. warneri* (5.6%) and one cluster each as *S. arlettae* (2.2%), *S. aureus* (2.2%), *S.*
367 *epidermidis* (1.1%), *S. haemolyticus* (1.1%), *S. hominis* (1.1%), *S. pasteurii* (1.1%), *S.*
368 *condiment* (1.1%), *S. piscifermentans* (1.1%), *S. saprophyticus* (1.1%), *S. simulans*
369 (18.9%), and *Macrococcus caseolyticus* (previously designated as *S. caseolyticus*,
370 5.5%). In general, *S. sciuri* was the most frequently isolated species and was dominant
371 in the samples of Bikalga (42.9%) and particularly Ntoba Mbodi (71.1%) where it
372 occurred from the beginning of the fermentation and persisted throughout the process.
373 In Soumbala, *S. simulans* (54.8%) was the dominant species. The isolate of *S.*
374 *haemolyticus* was clearly identified by 16S rRNA gene sequencing/EZtaxon analysis
375 with 100 % similarity, but the *rpoB* gene sequencing generated a 96.5% (30 base pairs
376 difference) similarity with *S. hominis* and 93.5% (30 base pairs difference) with *S.*
377 *haemolyticus*. The *spa* type of the *S. aureus* isolate was identified as t355.

378 The species of *S. sciuri* and *S. gallinarum* were common to the two countries.
379 One cluster (5) of *S. sciuri* was detected in all three types of food although this
380 particular cluster was not present in all samples (Table 3). Clusters 4 and 6 of the
381 same species were detected in samples from both countries but not in all three
382 products. *Staphylococcus gallinarum* was also recovered from both countries but not
383 all products, with one cluster (15) observed in Soumbala and Ntoba Mbodi. Nine
384 species were specific to Burkina Faso including five isolated solely from Soumbala (*S.*
385 *epidermidis*, *S. pasteurii*, *S. condimentii*, *S. piscifermentans*, *S. simulans*) and four from
386 Bikalga (*S. aureus*, *S. cohnii*, *S. haemolyticus*, *M. caseolyticus*). On the other hand,
387 four species were peculiar to Congo/ Ntoba Mbodi (*S. arlettae*, *S. hominis*, *S.*
388 *saprophyticus*, *S. warneri*). Moreover, two clusters of *S. gallinarum* (1, 23) and two of
389 *S. sciuri* (2, 7) were peculiar to Soumbala while one cluster of *S. sciuri* was particular
390 to Bikalga.

391 3.2 Haemolysis on blood agar and toxin production

392 Haemolytic activity of the isolates on sheep blood agar was variable as seen in
393 Table 4. Positive haemolysis was observed mainly in isolates of *S. aureus* and *S.*
394 *simulans*. *Staphylococcus epidermidis*, *S. haemolyticus*, *S. pasteurii* and *S. warneri*
395 exhibited partial haemolysis, whereas the rest of the species did not show haemolytic
396 activity within the 48 h of incubation. The different species clusters showed similar
397 haemolysis potential.

398 None of the bacteria screened produced the staphylococcal enterotoxins A, B,
399 C and D. Furthermore, the genes coding for the pre-cited enterotoxins as well as the
400 other eight genes were not detected in the isolates.

401 3.3 Antimicrobial resistance

402 All bacteria were susceptible to gentamycin, kanamycin, streptomycin, and
403 vancomycin. For the rest of the antimicrobials, susceptibility was variable according to
404 the species, the cluster, and the antimicrobial screened (Table 4). Most of the
405 staphylococci were resistant to trimethoprim with 87.8 % of the isolates, representing
406 11 species, exhibiting phenotypic resistance to that antimicrobial. Both *S. cohnii*
407 clusters exhibited the highest phenotypic resistance profiles involving 11
408 antimicrobials. The bacteria with the least phenotypic resistance were *S. condimenti*
409 and *S. pasteuri*, which showed reduced susceptibility toward sulfamethoxazole only.
410 Within the same species, the different clusters exhibited different resistance patterns,
411 but all isolates from the same cluster showed a similar pattern irrespective of the origin.
412 For example, isolates of cluster 1 of *S. gallinarum* were resistant to ceftiofur,
413 clindamycin, quinupristin/dalfopristin and tiamulin, whereas those of cluster 15 were
414 resistant to clindamycin, fusidate, penicillin, quinupristin/dalfopristin, tiamulin,
415 sulfamethoxazole, and trimethoprim, resulting in phenotypic resistance profiles which
416 differ by five antimicrobials (Table 4).

417 None of the 38 AMR genes screened was detected in the isolates of *S.*
418 *condimenti*, *S. pasteuri*, *S. piscifermentans*, *S. simulans*, and *M. caseolyticus* (Table
419 4). For the other species, different genes encoding resistance to multiple
420 antimicrobials including penicillin: *bla_ZA*, chloramphenicol: *cat501*, trimethoprim:
421 *dfr(A)*, *dfr(G)*, erythromycin: *msr(A)*, methicillin: *mecA*, *mecA1* and tetracycline: *tet (K)*
422 were detected, with a degree of AMR genotypic profile variability observed between
423 some species and clusters of the same species (Table 4). The *mecA1* gene was
424 detected in 45.6% of the isolates, *mecA* in 20%, *tet (K)* in 11.1 %, *bla_ZA* in 10 %, *dfr(A)*
425 in 10%, *msr(A)* in 6.7%, *cat501* in 1.1%, and *dfr(G)* in 1.1 %. Isolates of all six clusters
426 of *S. sciuri* showed *mecA1*, but bacteria in only three clusters displayed *mecA*. None

427 of the *S. aureus* isolates contained *mecA*, *mecA1* or *mecC*. The isolates of *S.*
428 *epidermidis* and *S. warneri* showed the highest numbers (4) of AMR genes, including
429 *blazA*, *dfr(A)*, *dfr(G)* and *tet(K)* for *S. epidermidis* and *blazA*, *Cat501*, *dfr(A)* and *msr(A)*
430 for cluster 13 of *S. warneri*. The *blazA*, *mecA*, *mecA1* and *tet (K)* genes were found in
431 isolates from all products and all countries, *cat501* in Ntoba Mbodi /Congo, *dfr(A)* in
432 Ntoba Mbodi/Congo and Soumbala/Burkina Faso, *dfr(G)* in Soumbala/Burkina Faso,
433 and *msr(A)* in Ntoba Mbodi /Congo and Bikalga/Burkina Faso

434 Conjugation experiments produced *E. faecalis* JH2-2 PETs with increased
435 resistance to erythromycin and tetracycline. The MIC of some PETs increased up to 6
436 and 4 fold for erythromycin and tetracycline respectively (Table 5). All PETs showed
437 the same rep-PCR profiles as that of the recipient *E. faecalis* JH2-2 (results not
438 shown). No bacteria were recovered on the antimicrobial selective agar plates
439 inoculated with cells which did not undergo a conjugation process. Positive amplicons
440 were obtained for the genes screened for transfer using both total and plasmid DNA
441 samples from the donors. However, no positive amplicon was observed with DNA
442 samples from the recipient and the PETs under the experimental conditions used.

443 **4. Discussion**

444 The specificity in the bacterial profile of the products can be attributed to
445 differences in the raw materials and production processes. Also, environmental factors
446 can affect bacterial populations in a given geographical area and its associated plants
447 and thus can affect the bacterial population of the products derived from such plants.
448 However, since a similar type of fermentation occurs during production of the
449 fermented foods, similarities can be anticipated. High bacterial counts as observed in
450 our study have been reported in other studies describing the microbial composition of
451 alkaline fermented products (Ahaotu et al., 2013; Azokpota et al., 2006; Dakwa et al.,

2005; Ouoba et al., 2010). Furthermore, the increase of pH constitutes a selection of analogous alkali resistant microorganisms in the foods. The presence of the same clusters of *S. sciuri* and *S. gallinarum* in both countries suggests a wide distribution of those strains in the environment. A peculiarity of the *S. sciuri* isolates recovered from the three products is that they were all oxidase negative. This seems unusual because the *S. sciuri* group of species including *S. sciuri*, *S. fleuritti*, *S. lentus*, *S. stepanovicii*, and *S. vitulinus* are known as the only oxidase positive species of the *Staphylococcus* genus (Becker et al., 2014). The discrepancy could be related to the origin specificity and to a lesser extent, the screening methods. The fact that the *S. aureus* PFGE profile did not match that of any clinical and environmental isolates in the CDC database suggests that the cluster is uncommon and not globally distributed.

The species of *S. sciuri*, *S. gallinarum*, *S. epidermidis*, *S. aureus* and *M. caseolyticus* have been reported in other African alkaline fermented products such maari, dawadawa, and soydawadawa (Parkouda et al., 2009). However, most of the other species, as well as the high species and strain diversity in the *Staphylococcus* profiles observed in the current study, seem to have not been reported before. The reason could be that *Staphylococcus* populations in such foods are usually screened as part of the general microbial population and are not addressed specifically and thoroughly using both phenotypic and genotypic methods. The species of *S. condimenti* and *S. piscifermentans* have been reported to be typically associated with fermented foods such as fermented fish, sausages and their starter cultures (Becker et al., 2014). It is believed that *Staphylococcus* spp. from fermented foods are contaminants from the raw materials, equipment, environment and possibly the producers. To produce Ntoba Mbodi, no heating step is included in the process, thus the original *Staphylococcus* isolates, derived from the cassava and wrapping leaves,

477 that can withstand alkaline conditions probably constitute the major part of the
478 staphylococcal population. However, production of Soumbala and Bikalga includes
479 long cooking steps (24-48h), so it can be assumed that there is a recontamination of
480 the products from the processing and storage materials, the producers and the
481 environment, especially during the drying process. The limited recovery of
482 *Staphylococcus* isolates (90/400 presumptive isolates) supports their status as a minor
483 component of the microflora of the foods. They have often been isolated from alkaline
484 fermented foods as secondary microorganisms, but a clear indication of their role in
485 the fermentation is not established. The main microorganisms responsible for alkaline
486 fermentation, and which bring about desired changes to the foods, have been
487 demonstrated to be *Bacillus* species (Odunfa and Adewuyi, 1985; Ouoba et al., 2017).
488 Fermentation trials conducted by Odunfa and Adewuyi (1985) showed that *S.*
489 *saprophyticus* was unable to ferment African locust bean seeds to generate a product
490 with the typical organoleptic and nutritional properties. Nevertheless, it can be
491 suggested that the presence of *Staphylococcus* species contributes to some
492 biochemical activities such as those related to the degradation of proteins and lipids
493 that some species are able to perform during meat fermentation to produce sausages
494 (Casaburi et al., 2008; Milicevic et al., 2014).

495 *Staphylococcus* spp., especially CoPS, are known to cause various illnesses
496 including food intoxication due to their capacity to produce heat resistant enterotoxins.
497 Interestingly, none of the isolates investigated, including the two CoP *S. aureus*,
498 exhibited a potential toxin-producing phenotype and genotype. It cannot be ruled out
499 that they may contain other toxin genes that were not screened, but since the most
500 common genes were not detected, it can be assumed that these bacteria do not
501 represent a serious threat for consumers. In Burkina Faso and the Republic of the

502 Congo, there are no official reports of foodborne illnesses resulting from the
503 consumption of alkaline traditional fermented foods, although this may be related to
504 non-availability of data and lack of statistics. If unrecorded cases had occurred, these
505 would probably have been attributed to toxins elaborated by the fermenting *B. cereus*
506 strains that commonly occur in the products (Ouoba et al., 2008 b; Vouidibio-Mbozo
507 et al., 2017). In the study by Fowoyo and Ogunbanwo (2017a), the *sea*, *seb*, *sec* and
508 *sed*, and *tsst-1* genes were detected in CoNS, including *S. epidermidis*, *S. simulans*,
509 *S. xyloso*, *S. kloosii* and *S. caprae*, isolated from six Nigerian traditional fermented
510 foods including iru, an alkaline fermented African locust bean product that is similar to
511 Soumbala. The positive amplicons were not sequenced to confirm the identity of the
512 gene, but the ability of some isolates to produce the staphylococcal toxins A, B, C, and
513 D suggest that these bacteria may pose a potential threat to consumers.

514 Antimicrobial resistances that are not associated with the presence of a gene
515 are of concern, but those involving a gene represent an enhanced threat, due to the
516 potential for transfer to other bacteria by mechanisms such as horizontal transfer, and
517 therefore can be spread. The *tet* (K), *blazA*, *dfr*(A), *msr*(A), *cat501*, and *dfr*(G) genes
518 detected can be associated with transferable mobile genetic elements such as
519 plasmids and transposons, and can therefore potentially be transferred to other
520 bacteria including pathogenic microorganisms. In the current research, *tet* (K), *blazA*,
521 *msr*(A), *cat501* and *dfr*(A) genes were amplified using plasmid DNA, suggesting that
522 they are located on plasmids, at least in the isolates screened. The AMR threat is
523 greater with bacteria with multi-drug genotypic profiles, such as one of the isolates of
524 *S. warneri* which exhibited AMR genes for four antimicrobials: erythromycin,
525 chloramphenicol, penicillin and trimethoprim, and those of *S. epidermidis* and *S.*

526 *hominis* which showed respectively four and three genes encoding resistance to
527 penicillin, tetracycline and chloramphenicol.

528 Methicillin resistance is an important factor when screening AMR in
529 *Staphylococcus*. This is because MRSA is responsible for a serious and persistent
530 problem in treatment of infections caused by such bacteria. The most virulent MRSA
531 are also resistant to other beta-lactam antimicrobials. Methicillin resistance is usually
532 caused by the presence of the *mecA* gene coding for a penicillin-binding protein
533 (PBP2a) associated with a significantly lower affinity for beta-lactams. Thus, cell-wall
534 biosynthesis by the bacteria continues, even when a typically inhibitory concentration
535 of antibiotic is administered (Peacock et al., 2015). Variants of *mecA* include genes
536 such as *mecA* and *mecC* (*mecA*_{LGA251}) originally found in *S. aureus* isolates, *mecA1*,
537 *mecA2* and *mecB* (*mecAm*) initially detected in the *S. sciuri* group, *S. vitulinus* and *M.*
538 *caseolyticus* respectively (Ito et al., 2012). Isolates of three clusters of *S. sciuri* from
539 both countries exhibited both *mecA1* and *mecA*, whereas the isolates of the three
540 other clusters also found in both countries exhibited only *mecA1*. The presence of the
541 *mecA1* gene in *S. sciuri* is a common trait of this species and does not generally imply
542 methicillin resistance in the bacteria (Couto et al., 2000). However, the presence of
543 *mecA* in the isolates implicates methicillin resistance and represents a threat, although
544 a phenotypic resistance was not observed for ceftiofur. Couto et al. (2000) showed
545 that only *S. sciuri* isolates that possess a copy of *mecA* in addition to *mecA1* exhibit
546 significant levels of resistance to beta-lactams. In our study, such isolates displayed
547 phenotypic resistance to penicillin. Interestingly, the *S. aureus* did not display any of
548 the methicillin resistance genes screened and were susceptible to ceftiofur. However,
549 the strains were resistant to penicillin and contained the *blaZ* gene.

550 The prevalence of AMR genes in various CoNS and CoPS from processed
551 and unprocessed foods such as meat, fish and fermented dairy and meat products
552 from different origins has been widely documented (Argudín et al., 2011; Kürekci,
553 2016; Perreten et al., 1998; Zdolec et al., 2012). However, such information on CoNS
554 and CoPS from alkaline fermented products and especially those from Africa are, to
555 the best of our knowledge, non-existent. This can be due to the limited importance
556 attributed to these bacteria in alkaline fermentation. The study by Fowoyo and
557 Ogunbanwo (2017b) on CoNS from different Nigerian fermented foods, reported
558 phenotypic resistance to antimicrobials such as trimethoprim-sulfamethoxazole,
559 ciprofloxacin, erythromycin, gentamicin, and cefotaxime, but no AMR gene was
560 reported, probably because only *mecA* was screened in the isolates with no positive
561 results.

562 The significant increase of the MIC above the cut-off values in the PETs for
563 erythromycin and tetracycline indicates that they have acquired resistance to the
564 antimicrobials. Since the genes screened were not detected in the PETs, it can be
565 assumed that they were not transferred, but that the isolates acquired resistance
566 through other mechanisms, such as mutations. The occurrence of resistances was
567 triggered by the presence of the *Staphylococcus* donors because PETs were not
568 recovered from recipient cells which did not undergo a conjugation process. Mutations
569 can occur as a result of a mechanism by which the *Staphylococcus* spp. stimulate an
570 evolution of *E. faecalis* JH2-2 by increasing, for instance, the expression of error-prone
571 DNA repair mechanisms (Naghizadeh Raeisi, 2018). Alternatively, the growth of the
572 recipient could be stimulated by the donor leading to a larger recipient population in
573 which mutations can arise (Naghizadeh Raeisi, 2018). However, it is worth noting that
574 when a low number of plasmid copies which carry a gene is transferred, it may not be

575 detected using conventional PCR. Moreover, other genes coding for resistance to the
576 antimicrobials and which were not screened, may be present in the donors and
577 transferred to the recipients. Transfer potential varies according to different factors
578 related to the environment, type of donor and recipient, and the location of the AMR
579 genes. Thus, the genes screened may have not been transferred into *E. faecalis* JH2-
580 2 and therefore the threat with regards to the potential of the donors to spread AMR is
581 limited. However, the possibility of gene transfer to other types of bacteria cannot be
582 ruled out since the genes are located on plasmids which are mobile genetic elements
583 that can be transferred in appropriate conditions.

584 The current study demonstrates ecological similarities and differences in the
585 *Staphylococcus* population of different types of alkaline fermented foods. The main
586 microorganisms identified are primarily CoNS with a very limited number of CoPS, as
587 would be expected in non-clinical samples. The bacteria did not suggest a safety issue
588 with regards to the potential to produce toxins responsible for foodborne diseases.
589 Due to the presence of multiple AMR genes in some isolates, they are reservoirs of
590 AMR determinants which may be spread under particular conditions. However, in
591 general, the results of the study herein reported suggest little safety importance of the
592 staphylococcus isolates in the fermented products screened.

593

594 **Acknowledgements**

595 The authors are grateful to the following people for their technical and scientific support
596 in the realization of the research project: Professor Simon C. Kobawila, and Professor
597 Delphin Louembe (Marien N’Gouabi University, Brazzaville, Congo), Dr Brigitte
598 Awamaria (London Metropolitan University, London, UK), Dr Rene S. Hendriksen,
599 Susanne Karlsmose Pedersen, Ana Rita Rebelo, and Dr. Lars Bøgo Jensen

600 (Technical University of Denmark, Lyngby, Denmark), Janne M. Benjaminsen
601 (University of Copenhagen, Copenhagen, Denmark), Dr Eva Møller Nielsen, and Dr
602 Jesper Larsen (Statens Serum Institut, Copenhagen, Denmark), Dr Molly Freeman,
603 Dr Jeniffer Concepcion Acevedo, and Valerie S. Albrecht (Centers for Disease Control
604 and Prevention, Atlanta, GA, USA)

605

606 **Declaration of interest:** None

607 **Funding:** This research did not receive any specific grant from funding agencies in
608 the public, commercial, or not-for-profit sectors

609 **References**

610 Ahaotu, I., Anyogu, A., Njoku, O.H., Odu, N.N., Sutherland, J.P., Ouoba, L.I.I., 2013
611 Molecular identification and safety of *Bacillus* species involved in the fermentation of
612 African oil beans (*Pentaclethra macrophylla* Benth) for production of Ugba. *Int. J. Food*
613 *Microbiol.* 162, 95-104.

614 Angulo, F., Nargund, V., Chiller, T., 2004. Evidence of an Association Between Use of
615 Anti-microbial Agents in Food Animals and Anti-microbial Resistance Among Bacteria
616 Isolated from Humans and the Human Health Consequences of Such Resistance. *J.*
617 *Vet. Med. B.* 51, 374-379.

618 Argudín, M., Tenhagen, B., Fetsch, A., Sachsenröder, J., Käsbohrer, A., Schroeter,
619 A., Hammerl, J.A., Hertwig, S., Helmuth, R., Bräunig, J., Mendoza, M.C., Appel, B.,
620 Rodicio, M.R., Guerra, B., 2011. Virulence and Resistance Determinants of German
621 *Staphylococcus aureus* ST398 Isolates from Non-human Sources. *Appl. Environ.*
622 *Microbiol.* 77, 3052-3060.

623 Azokpota, P., Hounhouigan, D.J., Nago, M.C., 2006. Microbiological and chemical
624 changes during the fermentation of African locust bean (*Parkia biglobosa*) to produce
625 *afitin*, *iru* and *sonru*, three traditional condiments produced in Benin. *Int. J. Food*
626 *Microbiol.* 107, 304 – 309.

627 Becker, K., Heilmann, C., Peters, G., 2014 Coagulase-Negative Staphylococci. *Clin.*
628 *Microbiol. Rev.* 27, 870-926.

629 Breckinridge, J., Bergdoll, M., 1971. Outbreak of Food-Borne Gastroenteritis Due to a
630 Coagulase-Negative Enterotoxin-Producing *Staphylococcus*. *N. Engl. J. Med.* 284,
631 541-543.

632 Casaburi, A., Di Monaco, R., Cavella, S., Toldrá, F., Ercolini, D., Villani, F., 2008.
633 Proteolytic and lipolytic starter cultures and their effect on traditional fermented
634 sausages ripening and sensory traits. *Food Microbiol.* 25, 335-347.

635 Couto, I., Santos Sanches, I., Sá-Leão, R., de Lencastre, H., 2000. Molecular
636 characterization of *Staphylococcus sciuri* strains isolated from humans. *J. Clin.*
637 *Microbiol.* 38, 136-143.

638 Crass, B., Bergdoll, M., 1986. Involvement of coagulase-negative staphylococci in
639 toxic shock syndrome. *J. Clin. Microbiol.* 23, 43-45.

640 Dakwa, S., Sakyi-Dawson, E., Diako, C., Annan, N.T., Amoa-Awua, W.K., (2005)
641 Effect of boiling and roasting on the fermentation of soybeans into dawadawa (soy-
642 dawadawa). *Int. J. Food Microbiol.* 104, 69 – 82.

643 Drancourt, M., Raoult, D., 2002. *rpoB* Gene Sequence-Based Identification of
644 *Staphylococcus* Species. *J. Clin. Microbiol.* 40, 1333-1338.

645 EUCAST., 2018. Breakpoint tables for interpretation of MICs and zone diameters.
646 http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_
647 [8.1_Breakpoint_Tables.pdf](http://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_8.1_Breakpoint_Tables.pdf)

648 EURL-AR., 2009. Protocol for MIC-testing with sensititre of *Staphylococcus* spp.
649 [https://www.eurl-ar.eu/CustomData/Files/Folders/21-protocols/285_1-protocol-for-](https://www.eurl-ar.eu/CustomData/Files/Folders/21-protocols/285_1-protocol-for-mic.pdf)
650 [mic.pdf](https://www.eurl-ar.eu/CustomData/Files/Folders/21-protocols/285_1-protocol-for-mic.pdf)

651 EURL-AR., 2013. List of primers for detection of antimicrobial resistance genes.
652 [https://www.eurl-ar.eu/CustomData/Files/Folders/25-resourcer/333_primerliste-til-](https://www.eurl-ar.eu/CustomData/Files/Folders/25-resourcer/333_primerliste-til-web-07-11-2013-9.pdf)
653 [web-07-11-2013-9.pdf](https://www.eurl-ar.eu/CustomData/Files/Folders/25-resourcer/333_primerliste-til-web-07-11-2013-9.pdf)

654 EURL-AR., 2017. PROTOCOL For antimicrobial susceptibility testing of *Escherichia*
655 *coli*, enterococci and staphylococci. [https://www.eurl-](https://www.eurl-ar.eu/CustomData/Files/Folders/23-eqas/356_protocol-eurl-ar-eqas-ec-ent-staph-2017.pdf)
656 [ar.eu/CustomData/Files/Folders/23-eqas/356_protocol-eurl-ar-eqas-ec-ent-staph-](https://www.eurl-ar.eu/CustomData/Files/Folders/23-eqas/356_protocol-eurl-ar-eqas-ec-ent-staph-2017.pdf)
657 [2017.pdf](https://www.eurl-ar.eu/CustomData/Files/Folders/23-eqas/356_protocol-eurl-ar-eqas-ec-ent-staph-2017.pdf).

658 EURL-AR., 2012. Protocol for amplification of *MecA*, *MecC* (MecALGA251), *spa* and
659 *PVL*. [https://www.eurl-ar.eu/CustomData/Files/Folders/21-protocols/279_pcr-spa-](https://www.eurl-ar.eu/CustomData/Files/Folders/21-protocols/279_pcr-spa-pvl-meca-mecc-sept12.pdf)
660 [pvl-meca-mecc-sept12.pdf](https://www.eurl-ar.eu/CustomData/Files/Folders/21-protocols/279_pcr-spa-pvl-meca-mecc-sept12.pdf)

661 EFSA., 2012. Technical specifications on the harmonised monitoring and reporting of
662 antimicrobial resistance in methicillin-resistant *Staphylococcus aureus* in food-
663 producing animals and food. EFSA J. 10, 2897.

664 Fonseca, S., Cachaldora, A., Gómez, M., Franco, I., Carballo, J., 2013. Monitoring the
665 bacterial population dynamics during the ripening of Galician chorizo, a traditional dry
666 fermented Spanish sausage. Food Microbiol. 33, 77-84.

667 Fowoyo, P., Ogunbanwo, S., 2017a. Virulence and toxigenicity of coagulase-negative
668 staphylococci in Nigerian traditional fermented foods. *Can. J. Microbiol.* 62, 572-578.

669 Fowoyo, P., Ogunbanwo, S., 2017b. Antimicrobial resistance in coagulase-negative
670 staphylococci from Nigerian traditional fermented foods. *Ann. Clin. Microbiol.*
671 *Antimicrob.* 16, 4.

672 Gregersen, T., 1978. Rapid method for distinction of gram-negative from gram-positive
673 bacteria. *Eu. J. Appl. Microbiol and Biotechnol.* 5, 123-127.

674 Gundogan, N., Citak, S., Yucel, N., Devren, A., 2005. A note on the incidence and
675 antibiotic resistance of *Staphylococcus aureus* isolated from meat and chicken
676 samples. *Meat Sci.* 69, 807-810.

677 Hennekinne, J., De Buyser, M., Dragacci, S., 2012. *Staphylococcus aureus* and its
678 food poisoning toxins: characterization and outbreak investigation. *FEMS Microbiol.*
679 *Rev.* 36, 815-836.

680 Yoon, S.H., Ha, S.M., Kwon, S., Lim, J., Kim, Y., Seo, H., Chun, J., 2017. Introducing
681 EzBioCloud: a taxonomically united database of 16S rRNA gene sequences and
682 whole-genome assemblies. *Int. J. Syst. Evol. Microbiol.* 67, 1613-1617.

683 Ito, T., Hiramatsu, K., Tomasz, A., de Lencastre, H., Perreten, V., Holden, M.,
684 Coleman, D.C., Goering, R., Giffard, P.M., Skov, R.L., Zhang, K., Westh, H., O'Brien,
685 F., Tenover, F.C., Oliveira, D.C., Boyle-Vavra, S., Laurent, F., Kearns, A.M.,
686 Kreiswirth, B., Ko, K.S., Grundmann, H, Sollid, J.E., John, J.F. Jr., Daum, R.,
687 Soderquist, B., Buist, G., IWG-SCC., 2012. Guidelines for Reporting Novel *mecA*
688 Gene Homologues. *Antimicrob. Agents Chemother.* 56, 4997-4999.

689 Johnson, W., Tyler, S., Ewan, E., Ashton, F., Pollard, D., Rozee, K., 1991. Detection
690 of Genes for Enterotoxins, Exfoliative Toxins, and Toxic Shock Syndrome Toxin 1 in
691 *Staphylococcus aureus* by the Polymerase Chain Reaction. J. Clin. Microbiol. 29, 426-
692 430.

693 Kiers, J.L, Van laeken, A.E., Rombouts, F.M., Nout, M.J., 2000. In vitro digestibility of
694 Bacillus fermented soya bean. In vitro digestibility of Bacillus fermented soya bean.
695 Int. J. Food Microbiol., 60,163-169.

696 Kurekci, C., Onen, S., Yipel, M., Aslantas, O., Gundogdu, A., 2016. Characterisation
697 of phenotypic and genotypic antibiotic resistance profile of enterococci from cheeses
698 in Turkey. Korean J. Food Sci. Anim. Resour. 36, 352-358.

699 LPSN., 2018 - List of prokaryotic names with standing in nomenclature [Internet].
700 Bacterio.net. <http://www.bacterio.net/>

701 Milicevi, B., Danilovic, B., Zdolec, N., Kozachinski, L., Dobranic, V., Savic, D., 2014.
702 Microbiota of the fermented sausages: influence to product quality and safety. Bulg. J.
703 Agric Sci. 20, 1061-1078.

704 Naghizadeh Raeisi, S., Ghodduzi, H., Juncker Boll, E., Farahmand, N., Stuer-
705 Lauridsen, B., Johansen, E., Sutherland, J.P., Ouoba, L.I.I., 2018. Antimicrobial
706 susceptibility of bifidobacteria from probiotic milk products and determination of the
707 genetic basis of tetracycline resistance in *Enterococcus* species after in vitro
708 conjugation with *Bifidobacterium animalis subsp. lactis*. Food Control. 2018; 94: 205-
709 211.

710 Nashev, D., Toshkova, K., Isrina, S., Salasia, S., Hassan, A., Lammler, C., Zschock,
711 M., 2004. Distribution of virulence genes of *Staphylococcus aureus* isolated from
712 stable nasal carriers. FEMS Microbiol. Lett. 233, 45-52.

713 Odunfa, S., Adewuyi, E., 1985. Optimization of process conditions for the fermentation
714 of African locust bean (*Parkia biglobosa*) (Effect of starter cultures). Chem. Mikrobio.
715 Technol. Lebensm. 9, 6-10.

716 Omoe, K., Ishikawa, M., Shimoda, Y., Hu, D., Ueda, S., Shinagawa, K., 2002.
717 Detection of seg, seh, and sei genes in *Staphylococcus aureus* isolates and
718 determination of the enterotoxin productivities of *S. aureus* isolates harboring seg, seh,
719 or sei genes. J. Clin. Microbiol. 40: 857-862.

720 Ouoba, L., Parkouda, C., Diawara, B., Scotti, C., Varnam, A., 2008a. Identification of
721 *Bacillus* spp. from Bikalga, fermented seeds of *Hibiscus sabdariffa*: phenotypic and
722 genotypic characterization. J. Appl. Microbiol. 104, 122-131.

723 Ouoba, L., Thorsen, L., Varnam, A., 2008b. Enterotoxins and emetic toxins production
724 by *Bacillus cereus* and other species of *Bacillus* isolated from Soumbala and Bikalga,
725 African alkaline fermented food condiments. Int. J. Food Microbiol. 124, 224-230.

726 Ouoba, L., Lei, V., Jensen, L., 2008c. Resistance of potential probiotic lactic acid
727 bacteria and bifidobacteria of African and European origin to antimicrobials:
728 Determination and transferability of the resistance genes to other bacteria. Int. J. Food
729 Microbiol. 121, 217-224.

730 Ouoba, L., Nyanga-Koumou, C.A., Parkouda, C., Sawadogo, H., Kobawila, S.C.,
731 Keleke, S., Diawara, B., Louembe, D., Sutherland, J.P., 2010. Genotypic diversity of

732 lactic acid bacteria isolated from African traditional alkaline-fermented foods. Int. J.
733 Food Microbiol. 108, 2019 – 2029.

734 Ouoba, L., 2015. Bikalga/Furundu/Mbuja. In: Sarkar, P., Nout, M.J.R., (Eds.),
735 Handbook of indigenous foods involving alkaline fermentation. CRC Press, Boca
736 Raton, pp. 129-136.

737 Ouoba, L., 2017. Traditional alkaline fermented foods: Selection of functional *Bacillus*
738 starter cultures for Soumbala production. In: Speranza, B., Bevilacqua, A., Corbo, M.,
739 Sinigaglia, M., (Eds), Starter Cultures in Food Production. 1st ed. Wiley Blackwell,
740 Chichester, pp. 370-383.

741 Parkouda, C., Diawara, B., Ouoba, L., 2008. Technology and physico-chemical
742 characteristics of Bikalga, alkaline fermented seeds of *Hibiscus sabdariffa*. Afr. J.
743 Biotechnol. 7, 916-922.

744 Parkouda, C., Nielsen, D., Azokpota, P., Ouoba, L.I.I., Amoa-Awua, W., Thorsen, L.,
745 Hounhouigan., J.D., Jensen, J.S, Tano-Debrah, K., Diawara, B., Jakobsen, M., 2009.
746 The microbiology of alkaline-fermentation of indigenous seeds used as food
747 condiments in Africa and Asia. Crit. Rev. Microbiol. 35,139-156.

748 Peacock, S., Paterson, G., 2015. Mechanisms of methicillin resistance in
749 *Staphylococcus aureus*. Annu. Rev. Biochem. 84, 577-601.

750 Perreten, V., Giampà, N., Schuler-Schmid, U., Teuber, M., 1998. Antibiotic resistance
751 genes in coagulase-negative staphylococci isolated from food. Syst. Appl. Microbiol.
752 21,113-120.

753 Podkowik, M., Park, J., Seo, K., Bystroń, J., Bania, J., 2013. Enterotoxigenic potential
754 of coagulase-negative staphylococci. Int. J. Food Microbiol. 163, 34-40.

755 Simeoni, D., Rizzotti, L., Cocconcelli, P., Gazzola, S., Dellaglio, F., Torriani, S., 2008.
756 Antibiotic resistance genes and identification of staphylococci collected from the
757 production chain of swine meat commodities. *Food Microbiol.* 25,196-201.

758 Soares Casaes Nunes, R., Mere Del Aguila, E., Paschoalin, V., 2015. Safety
759 Evaluation of the coagulase-negative staphylococci microbiota of salami:
760 superantigenic toxin production and antimicrobial resistance. *BioMed Res. Int.* 2015,
761 1-17.

762 Stegger, M., Andersen, P., Kearns, A., Pichon, B., Holmes, M., Edwards, G., Laurent,
763 F., Teale, C., Skov, R., Larsen, A.R., 2012. Rapid detection, differentiation and typing
764 of methicillin-resistant *Staphylococcus aureus* harbouring either *mecA* or the new
765 *mecA* homologue *mecALGA251*. *Clin. Microbiol. Infect.* 18, 395-400.

766 Sutcliffe, J., Grebe, T., Tait-Kamradt, A., Wondrack, L., 1996. Detection of
767 erythromycin-resistant determinants by PCR. *Antimicrob. Agents Chemother.* 40,
768 2562-2566.

769 Tsubakishita, S., Kuwahara-Arai, K., Sasaki, T., Hiramatsu, K., 2010. Origin and
770 Molecular evolution of the determinant of methicillin resistance in staphylococci.
771 *Antimicrob. Agents Chemother.* 10, 4352-4359

772 van Balen, J., Kelley, C., Nava-Hoet, R., Bateman, S., Hillier, A., Dyce, J., Wittum,
773 T.E., Hoet, A. E. 2013. Presence, Distribution, and Molecular Epidemiology of
774 Methicillin-Resistant *Staphylococcus aureus* in a Small Animal Teaching Hospital: A
775 Year-Long Active Surveillance Targeting Dogs and Their Environment. *Vector Borne*
776 *Zoonotic Dis.* 5, 299-311.

777 Viridis, S., Scarano, C., Cossu, F., Spanu, V., Spanu, C., De Santi, E. 2010. Antibiotic
778 resistance in *Staphylococcus aureus* and Coagulase Negative Staphylococci Isolated
779 from Goats with Subclinical Mastitis. *Vet. Med. Int.*, 2010, 517060.

780 Voudibio Mbozo, B. A., Kobawila, S., Anyogu, A., Awamaria, B., Louembe, D.,
781 Sutherland, J., Ouoba, L.I.I., 2017. Investigation of the diversity and safety of the
782 predominant *Bacillus pumilus sensu lato* and other *Bacillus* species involved in the
783 alkaline fermentation of cassava leaves for the production of Ntoba Mbodi. *Food*
784 *Control.* 82, 154-162.

785 Zdolec, N., Dobranić, V., Zdolec, G., Đuričić, D. 2013. Antimicrobial resistance of
786 coagulase-negative staphylococci and lactic acid bacteria from industrially produced
787 dairy products. *Mljekarstvo.* 63, 30-35.

788

Accepted Manuscript

789 **Figure caption**

790 Figure 1: Cluster analysis (Bionumerics, Dice's Coefficient of similarity, Unweighted
791 pair group method with arithmetic: UPGMA) of the different rep-PCR and PFGE
792 fingerprints of *Staphylococcus species* isolated from Bikalga, Soumbala and Ntoba
793 Mbodi

794

795

796

797

798

799

800

801

802

803

804

805

806

807

808

809

Accepted Manuscript

810 Table 1: Antimicrobials studied and their effect on the *Staphylococcus* isolates
 811 screened

Antimicrobials	Concentration Range screened (mg/L)	Cut off values(mg/L)	Genes screened ^e	<i>Staphylococcus</i> isolates		
				MIC Range (mg/L)	Resistant (%)	Susceptible (%)
Cefoxitin	0.5 - 16	4 ^{a,b}	<i>mecA</i> , <i>mecA1</i> and <i>mecC</i> for methicillin resistance	1 - 8	8.9	91.1
Chloramphenicol	4 - 64	8 ^c	<i>cat501</i> , <i>catA1</i> , <i>cmIA</i>	<4 - 64	52.2	47.8
Ciprofloxacin	0.25 - 8	1 ^a		<0.25 - 2	1.1	98.9
Clindamycin	0.12 - 4	0.25 ^a		<0.12 - 4	41.1	58.9
Erythromycin	0.25 - 8	1 ^a	<i>erm(A)</i> , <i>erm(B)</i> , <i>erm(C)</i> , <i>erA</i> , <i>erB</i> , <i>msrA</i> , <i>mphA</i> , <i>mefA/mefE</i>	<0.25 - >8	11.1	88.9
Fusidate	0.5 - 4	1 ^c		<0.5 - >4	43.3	56.7
Gentamicin	1 - 16	2 ^a	<i>aac(6')aph(2'')</i> , <i>aac(3'')II</i> , <i>aac(3'')IV</i> , <i>ant(2'')-I</i>	<1	0	100
Kanamycin	4 - 64	16 ^d		<4	0	100
Linezolid	1 - 8	4 ^a		<1 - 8R	1.1	98.9
Mupirocin	0.5 - 256	1 ^a		<0.5 - > 256	1.1	98.9
Penicillin	0.12 - 2	0.125 ^b	<i>BlaZ</i>	<0.12 - >2	43.3	56.7
Quinupristin-dalfopristin	0.5 - 4	1 ^a	<i>VatD</i> , <i>VatE</i>	<0.5 - 4	64.4	35.6
Rifampicin	0.016 - 0.5	0.5 ^c		<0.016 - >0.5	4.4	95.6
Streptomycin	4 - 32	16 ^b	<i>StrA</i> , <i>StrB</i> , <i>aadA</i> , <i>aadE</i>	<4 - 16	0	100
Sulfamethoxazole	64 - 512	128 ^a		<64 - >512	61.1	38.9
Tetracycline	0.5 - 16	1 ^a	<i>tet(M)</i> , <i>tet(L)</i> , <i>tet(S)</i> , <i>tet(Q)</i> , <i>tet(K)</i> , <i>tet(O)</i> , <i>tet(W)</i>	<0.5 - >16	6.7	93.76
Tiamulin	0.5 - 4	2 ^a		<0.5 - > 4	48.9	51.1
Trimethoprim	2 - 32	2 ^a	<i>dfr(A)</i> , <i>dfr(G)</i> , <i>dfr(K)</i>	<2 - >32	87.8	12.2
Vancomycin	1 - 16	2 ^a	<i>Van A</i> , <i>VanB</i> , <i>VanX</i>	<1	0	100

812

813 Cut-off values according to a) EFSA (2012), b) *EUURL-AR (2017)*, c) EUCAST (2018) and d) Virdis et
 814 al. (2010)

815 e) Most primer sequences are from *EUURL-AR (2013)* and Ouoba et al. (2008), except for *mecA1*
 816 (*Tsubakishita et al., 2010*), *erA*, *erB*, *msrA*, *mphA*, *mefA/mefE* (*Sutcliffe et al., 1996*), *dfr(G)*, *dfr(K)*
 817 (*Argudín et al., 2011*)

818 f) MIC: Minimal Inhibitory Concentration

819

820

821

822

823

824 Table 2: Identity and origin of the *Staphylococcus* species recovered from Soumbala,
 825 Bikalga, and Ntoba Mbodi

Samples/Origin	Fermentation time/ sampling site ^a	Isolates	^b Rep-PCR/ PFGE clusters	Identification by 16S RNA and <i>rpoB</i> gene sequencing		
Soumbala						
Burkina Faso						
S3	Site 1/ 0h	S35M (47)	14	<i>Staphylococcus epidermidis</i>		
S5	Site 1/ 48h	S51B (33)	23	<i>Staphylococcus gallinarum</i>		
		S54a M (34)	23	<i>Staphylococcus gallinarum</i>		
		S54b M (35)	23	<i>Staphylococcus gallinarum</i>		
		S63aM (41)	3	<i>Staphylococcus simulans</i>		
S6	Site 1/ Dried	SF1M (1)	1	<i>Staphylococcus gallinarum</i>		
SF	Site 2/ ≥72h	SF3M (2)	3	<i>Staphylococcus simulans</i>		
		SF2M (3)	5	<i>Staphylococcus sciuri</i>		
		SF2bB (6)	3	<i>Staphylococcus simulans</i>		
		SF2aB (7)	3	<i>Staphylococcus simulans</i>		
		SF1B (8)	3	<i>Staphylococcus simulans</i>		
		SF8M (21)	2	<i>Staphylococcus sciuri</i>		
		SF7M (22)	3	<i>Staphylococcus simulans</i>		
		SF6M (23)	7	<i>Staphylococcus sciuri</i>		
		SP6M (26)	10	<i>Staphylococcus piscifermentans</i>		
		SP5M (27)	3	<i>Staphylococcus simulans</i>		
		SP3M (28)	3	<i>Staphylococcus simulans</i>		
		SP1M (29)	11	<i>Staphylococcus pasteurii</i>		
		SP	Site 3/ ≥72h	SP4B (30)	3	<i>Staphylococcus simulans</i>
SP2B (31)	3			<i>Staphylococcus simulans</i>		
SP1B (32)	3			<i>Staphylococcus simulans</i>		
SZ	Site 4/ ≥72h			SZ1B (10)	3	<i>Staphylococcus simulans</i>
				SZ2B (11)	3	<i>Staphylococcus simulans</i>
				SZ6B (13)	3	<i>Staphylococcus simulans</i>
				SZ1M (14)	5	<i>Staphylococcus sciuri</i>
		SZ3M (15)	3	<i>Staphylococcus simulans</i>		
		SZ4M (16)	5	<i>Staphylococcus sciuri</i>		
		SZ5M (17)	9	<i>Staphylococcus condimentii</i>		
		SZ6M (18)	15	<i>Staphylococcus gallinarum</i>		
SZ7M (24)	3	<i>Staphylococcus simulans</i>				
SZ9M (25)	3	<i>Staphylococcus simulans</i>				
Bikalga						
Burkina Faso						
B2	Site 5/ 48h	B21M (50)	6	<i>Staphylococcus sciuri</i>		
		B22M (51)	4	<i>Staphylococcus sciuri</i>		
		B27aM (53)	16	<i>Staphylococcus cohnii</i>		
		B27bM (54)	16	<i>Staphylococcus cohnii</i>		
		B25M (56)	16	<i>Staphylococcus cohnii</i>		
		B25B (57)	8	<i>Staphylococcus sciuri</i>		
		B24B (58)	24	<i>Staphylococcus cohnii</i>		
		B3	Site 5/ 72h	B35M (60)	5	<i>Staphylococcus sciuri</i>
B36B (61)	5			<i>Staphylococcus sciuri</i>		
B34M (62)	5			<i>Staphylococcus sciuri</i>		
B32B (64)	5			<i>Staphylococcus sciuri</i>		
B31M (65)	5			<i>Staphylococcus sciuri</i>		
B31B (66)	8			<i>Staphylococcus sciuri</i>		

826 ^aSite 1: Ouagadougou (Samandin production site), Site 2: Fada market, Site 3: Ouagadougou (Patte
 827 d'Oie market), Site 4: Ouagadougou (Zogona market), Site 5: Ouagadougou (Zone Pilote production

828 site) ^b Rep-PCR : Repetitive element palindromic-polymerase chain reaction ; PFGE : Pulse Field Gel
829 Electrophoresis

Accepted Manuscript

830 Table 2 *continued*: Identity of the *Staphylococcus* species recovered from Soumbala,
 831 Bikalga, and Ntoba Mbodi

Samples/Origin	Fermentation time/ sampling site ^a	Isolates	Rep-PCR/ PFGE clusters	Identification 16S RNA and <i>rpoB</i> gene sequencing
B5	Site 5/ Dried	B57B (69)	18	<i>Staphylococcus aureus</i>
		B51B (70)	18	<i>Staphylococcus aureus</i>
BP	Site 6/ Dried	BP2B (73)	19	<i>Macrococcus caseolyticus</i>
		BP2M (74)	19	<i>Macrococcus caseolyticus</i>
		BP3B (75)	19	<i>Macrococcus caseolyticus</i>
		BP6B (76)	19	<i>Macrococcus caseolyticus</i>
		BP5B (77)	19	<i>Macrococcus caseolyticus</i>
BB Ntoba Mbodi	Site 7/ Dried	BB3B (84)	20	<i>S. haemolyticus</i>
Congo AU0	Site 8/ 0h	A04M (93)	25	<i>Staphylococcus sciuri</i>
		A05M (94)	12	<i>Staphylococcus warneri</i>
		A08M (95)	12	<i>Staphylococcus warneri</i>
		A07M (185)	12	<i>Staphylococcus warneri</i>
		A06B (193)	12	<i>Staphylococcus warneri</i>
AU1	Site 8/ 24h	A12B (96)	6	<i>Staphylococcus sciuri</i>
		A15M (104)	25	<i>Staphylococcus sciuri</i>
		A17M (105)	13	<i>Staphylococcus warneri</i>
		A18M (106)	17	<i>Staphylococcus arlettae</i>
		A15M (176)	4	<i>Staphylococcus sciuri</i>
		A16M (177)	6	<i>Staphylococcus sciuri</i>
		A111B (186)	5	<i>Staphylococcus sciuri</i>
		A110B (194)	5	<i>Staphylococcus sciuri</i>
AU2	Site 8/ 48h	A25B (111)	17	<i>Staphylococcus arlettae</i>
		A215B (113)	15	<i>Staphylococcus gallinarum</i>
		A21M (115)	4	<i>Staphylococcus sciuri</i>
		A27B (178)	5	<i>Staphylococcus sciuri</i>
		A29B (179)	6	<i>Staphylococcus sciuri</i>
		A26M (180)	6	<i>Staphylococcus sciuri</i>
		A23M (191)	4	<i>Staphylococcus sciuri</i>
AU3	Site 8/ 72h	A24M (192)	4	<i>Staphylococcus sciuri</i>
		A37bB (123)	6	<i>Staphylococcus sciuri</i>
		A39B (124)	21	<i>Staphylococcus hominis</i>
		A34M (128)	6	<i>Staphylococcus sciuri</i>
		A31M (188)	6	<i>Staphylococcus sciuri</i>
JO0	Site 9/ 0h	A33M (187)	6	<i>Staphylococcus sciuri</i>
		J03B (131)	6	<i>Staphylococcus sciuri</i>
		J04B (132)	6	<i>Staphylococcus sciuri</i>
		J01B (189)	4	<i>Staphylococcus sciuri</i>
		J02B (190)	6	<i>Staphylococcus sciuri</i>
JO3	Site 9/ 72h	J03M (138)	22	<i>Staphylococcus saprophyticus</i>
		J311B (148)	15	<i>Staphylococcus gallinarum</i>
		J33M (150)	6	<i>Staphylococcus sciuri</i>
		J34M (151)	6	<i>Staphylococcus sciuri</i>
		J31M (181)	6	<i>Staphylococcus sciuri</i>
MB	Site 10/ 72-96h	J32M (182)	6	<i>Staphylococcus sciuri</i>
		MB1M (156)	5	<i>Staphylococcus sciuri</i>
		MB11M (184)	5	<i>Staphylococcus sciuri</i>

832 ^aSite 5: Ouagadougou (Zone Pilote production site), Site 6: Ouagadougou (Patte d'Oie market), Site
 833 7: Bobo Dioulasso (market), Site 8: Brazzaville (Production site A), Site 9: Brazzaville (Production site
 834 B), Site 10: Brazzaville (market)

835 ^b Rep-PCR : Repetitive element palindromic-polymerase chain reaction ; PFGE : Pulse Field Gel
836 Electrophoresis

837

838

839

840

841

842

843

844

845

846

847

848

849

850

851

852

853

854

855

856

Accepted Manuscript

857 Table 3 Geographical distribution of the *Staphylococcus* species and associated clusters
 858

Species	Rep-PCR/ PFGE Clusters	Burkina Faso/ Soumbala				Burkina Faso/ Bikalga			Congo/ Ntoba Mbodi		
		Site 1 ^b	Site 2	Site 3	Site 4	Site 5	Site 6	Site 7	Site 8	Site 9	Site 10
<i>S. arlettae</i>	17	- ^c	-	-	-	-	-	-	+	-	-
<i>S. aureus</i>	18	-	-	-	-	+	-	-	-	-	-
<i>S. cohnii</i>	16	-	-	-	-	-	+	-	-	-	-
	24	-	-	-	-	-	+	-	-	-	-
<i>S. epidermidis</i>	14	+	-	-	-	-	-	-	-	-	-
<i>S. gallinarum</i>	1	+	-	-	-	-	-	-	-	-	-
	15	-	-	-	+	-	-	-	+	+	-
	23	+	-	-	-	-	-	-	-	-	-
<i>S. haemolyticus</i>	20	-	-	-	-	-	-	+	-	-	-
<i>S. hominis</i>	21	-	-	-	-	-	-	-	+	-	-
<i>S. pasteurii</i>	11	-	-	+	-	-	-	-	-	-	-
<i>S. condimentii</i>	9	-	-	-	+	-	-	-	-	-	-
<i>S. piscifermentans</i>	10	-	+	-	-	-	-	-	-	-	-
<i>S. saprophyticus</i>	22	-	-	-	-	-	-	-	-	+	-
<i>S. sciuri</i>	2	-	+	-	-	-	-	-	-	-	-
	4	-	-	-	-	+	-	-	+	+	-
	5	-	+	-	+	+	-	-	-	-	+
	6	-	-	-	-	+	-	-	+	+	-
	7	-	+	-	-	-	-	-	-	-	-
	8	-	-	-	-	+	-	-	-	-	-
	25	-	-	-	-	-	-	-	+	-	-
<i>S. simulans</i>	3	+	+	+	+	-	-	-	-	-	-
<i>S. warneri</i>	12	-	-	-	-	-	-	-	+	-	-
	13	-	-	-	-	-	-	-	+	-	-
<i>M. caseolyticus</i>	19	-	-	-	-	-	+	-	-	-	-

859
 860 ^a Rep-PCR : Repetitive element palindromic-polymerase chain reaction ; PFGE : Pulse Field Gel Electrophoresis

861 ^bSite 1: Ouagadougou (Samandin production site), Site 2: Fada market, Site 3: Ouagadougou (Patte d'Oie market), Site 4: Ouagadougou (Zogona market),
 862 Site 5: Ouagadougou (Zone Pilote production site), Site 6: Ouagadougou (Patte d'Oie market), Site 7: Bobo Dioulasso (market), Site 8: Brazzaville
 863 (Production site A), Site 9: Brazzaville (Production site B), Site 10: Brazzaville (market)

864 ^c - : not detected, + : detected

865 Table 4 Profiles of antimicrobial resistance (AMR) and toxin production potential of the species and clusters of *Staphylococcus* from
 866 Bikalga, Soumbala, and Ntoba Mbodi

Origin ^a	Rep-PCR/ PFGE Cluster	Species	Number of Isolates	AMR phenotype ^b	AMR genes	Toxin genes/ Toxin A, B, C, D	Haemolysis on blood agar
Ntoba Mbodi (Co)	17	<i>S. arlettae</i>	2	CHL, CLI, FUS, SYN, SMX, TIA, TMP	dfr(A)	- ^c	-
Bikalga (BF)	18	<i>S. aureus</i>	2	CHL, ERY, PEN, SYN, SUL	blazA	-	+
Bikalga (BF)	16	<i>S. cohnii</i>	3	FOX, CHL, CLI, ERY, FUS, PEN, RIF, SMX, SYN, TIA, TMP	msr(A)	-	-
Bikalga (BF)	24	<i>S. cohnii</i>	1	FOX, CLI, ERY, FUS, PEN, RIF, SMX., SYN, TET, TIA, TMP	msr(A)	-	-
Soumbala (BF)	14	<i>S. epidermidis</i>	1	CHL, PEN, SMX, TET, TMP	blazA, dfr(A), dfr(G) tet(K)	-	±
Soumbala (BF)	1	<i>S. gallinarum</i>	1	FOX, CLI, SYN, TIA	tet(K)	-	-
Soumbala, Ntoba Mbodi (BF, Co)	15	<i>S. gallinarum</i>	3	CLI, FUS, PEN, SYN, TIA, SMX, TMP	tet(K)	-	-
Soumbala (BF)	23	<i>S. gallinarum</i>	3	FOX, CLI, FUS, PEN, SYN, TIA, TMP	tet(K)	-	-
Bikalga (BF)	20	<i>S. haemolyticus</i>	1	CLI, ERY, PEN, SYN, TIA, TMP	tet(K), msr(A)	-	±
Ntoba Mbodi (Co)	21	<i>S. hominis</i>	1	CHL, PEN, SMX, TET, TMP	blazA, tet(K), dfr(A)	-	-
Soumbala (BF)	11	<i>S. pasteurii</i>	1	SMX		-	±
Soumbala (BF)	10	<i>S. piscifermentans</i>	1	CHL, ERY, SMX, TIA		-	-
Soumbala (BF)	9	<i>S. condimentii</i>	1	SMX		-	-
Ntoba Mbodi (Co)	22	<i>S. saprophyticus</i>	1	CHL, PEN, TET, TMP	tet(K)	-	-
Soumbala (BF)	2	<i>S. sciuri</i>	1	CLI, FUS, SMX, TIA, TMP	mecA, mecA1	-	-
Bikalga, Ntoba Mbodi (BF, Co)	4	<i>S. sciuri</i>	6	CLI, FUS, TIA, TMP	mecA1	-	-
Bikalga, Soumbala, Ntoba Mbodi (BF, Co)	5	<i>S. sciuri</i>	13	CLI, FUS, TMP	mecA1	-	-
Bikalga, Ntoba Mbodi (BF, Co)	6	<i>S. sciuri</i>	16	CHL, PEN, SYN, SMX, TIA, TMP	mecA, mecA1	-	-
Soumbala (BF)	7	<i>S. sciuri</i>	1	CLI, ERY, FUS, SYN, TIA, TMP	mecA, mecA1	-	-
Bikalga (BF)	8	<i>S. sciuri</i>	2	CHL, PEN, SYN, TET, TIA, TMP	mecA1	-	-
Ntoba Mbodi (Co)	25	<i>S. sciuri</i>	2	CLI, FUS, SMX, SYN, TIA, TMP	mecA1	-	-
Soumbala (BF)	3	<i>S. simulans</i>	17	CHL, LZD, SYN, SMX, TMP		-	+
Ntoba Mbodi (Co)	12	<i>S. warneri</i>	4	FUS, PEN, SYN, SMX, TMP	blazA, dfr(A)	-	±
Ntoba Mbodi (Co)	13	<i>S. warneri</i>	1	CHL, ERY, PEN, SMX, TMP	blazA, Cat501 dfr(A), msr(A)	-	-
Bikalga (BF)	19	<i>M. caseolyticus</i>	5	CIP, MUP		-	-

867 ^a BF: Burkina Faso, Co: Congo


868 ^b The profile represents the antimicrobials to which the isolates were resistant to; Cefoxitin: FOX, Chloramphenicol: CHL, Ciprofloxacin: CIP,
 869 Clindamycin: CLI, Erythromycin: ERY, Fusidate: FUS, Mupirocin: MUP, Quinupristin/Dalfopristin (Synercid): SYN, Linezolid: LZD, Penicillin:
 870 PEN, Rifampicin: RIF, Sulfamethoxazole: SMX, Tiamulin: TIA, Trimethoprim: TMP ^c -: negative, ±: partial haemolysis, +: positive haemolysis

871 Table 5: Minimal Inhibitory Concentration (MIC) for donors, recipient and potential *E. faecalis* JH2-2 transconjugants (PET)

MIC (mg/L): Erythromycin		MIC (mg/L): Tetracycline	
Donor <i>S. warneri</i>	>32	Donor <i>S. saprophyticus</i>	64
Recipient <i>E. faecalis</i>	<1	Recipient <i>E. faecalis</i>	1
PET-E1	16	PET-T1	8
PET-E2	<1	PET-T2	8
PET-E3	>32	PET-T3	16
PET-E4	<1	PET-T4	8
PET-E5	<1	PET-T5	8
PET-E6	4	PET-T6	16
PET-E7	<1	PET-T7	16
PET-E8	<1	PET-T8	4
PET-E9	>32	PET-T9	16
PET-E10	<1	PET-T10	4
PET-E11	32		
PET-E12	<1		
PET-E13	>32		
PET-E14	<1		

872

873



874

875 Figure 1

ACCEPT