

WestminsterResearch

<http://www.westminster.ac.uk/westminsterresearch>

Modification of bacterial cell membrane to accelerate decolorization of textile wastewater effluent using microbial fuel cells: role of gamma radiation, salinity and endogenous biosurfactant induction

Selim, N., Hamed, H., Fathy, R., Gamal, M., El Kareem, H.A., Breheny, M., Farahmand, N., Bowman, K., Kyazze, G., Keshavarz, T. and Gomaa, O.M.

A poster presented at the 4th European Meeting of the International Society for Microbial Electrochemistry and Technology, Newcastle University, Newcastle upon Tyne, 12 - 14 Sep 2018.

The WestminsterResearch online digital archive at the University of Westminster aims to make the research output of the University available to a wider audience. Copyright and Moral Rights remain with the authors and/or copyright owners.

Whilst further distribution of specific materials from within this archive is forbidden, you may freely distribute the URL of WestminsterResearch: (<http://westminsterresearch.wmin.ac.uk/>).

In case of abuse or copyright appearing without permission e-mail repository@westminster.ac.uk



Modification of bacterial cell membrane to accelerate decolorization of textile waste water effluent using Microbial Fuel Cell: role of gamma radiation, salinity and endogenous biosurfactant induction

Authors Nabila Selim¹, Heba Hamed², Reham Fathy², Marwa Gamal², Hussein Abd El Kareem², Kyle Bowman³, Mark Brehney³, Nasim Farhand³, Godfrey Kyazzae³, Tajalli Keshavarz³, Ola M. Gomaa^{2*}

Affiliation ¹Radiation Physics Department, ²Microbiology Department, National Center for Radiation Research and Technology (NCRRT), Egyptian Atomic Energy Authority (EAEA), Cairo-Egypt, ³Department of Life Sciences, University of Westminster, London, United Kingdom

Email*ola_gomaa@hotmail.com

Abstract

A combined approach was investigated to accelerate Microbial Fuel Cell (MFC) performance and textile wastewater decolorization through modifying bacterial membrane. The aim was to increase both bacterial adhesion on anode and electron mediator release. Ten Gram-positive exoelectrogenic bacteria were isolated from the anodic biofilm after decolorization of real textile waste water in mediator-less MFC. The isolates were identified and characterized, to understand the nature of the bacteria involved. According to the battery of tests performed, three factors gamma radiation, salinity and induction of endogenous biosurfactant were involved membrane modification. Dielectric measurement, a non-invasive technique, was used to measure the cell membrane permeability and cell surface charge. Plackett-Burman experimental design was carried out to determine the key contributor among the three studied factors. Exposing the cells to 1 KGy gamma radiation led to 7.84- and 1.71- fold increase in total surface-charge and cell-permeability, respectively. Scanning Electron Microscope (SEM) images and surface-bound protein concentrations for the samples indicated that biofilm formation increased under the same conditions. These results have been reflected on the power density profiles and decolorization of textile wastewater. Modification of bacterial membrane prior to MFC operation can be considered highly effective as a pre-treatment tool that accelerates MFC performance.