

WestminsterResearch

<http://www.westminster.ac.uk/westminsterresearch>

Effects of upper-body, lower-body, or combined resistance training on the ratio of follistatin and myostatin in middle-aged men

Bagheri, R., Rashidlamir, A., Motevalli, M., Elliott, B., Mehrabani, J. and Wong, A.

This is an author's accepted manuscript of an article published in the European Journal of Applied Physiology, pp. 1-11, 2019.

The final publication is available at Springer via:

<https://dx.doi.org/10.1007/s00421-019-04180-z>

The WestminsterResearch online digital archive at the University of Westminster aims to make the research output of the University available to a wider audience. Copyright and Moral Rights remain with the authors and/or copyright owners.

Whilst further distribution of specific materials from within this archive is forbidden, you may freely distribute the URL of WestminsterResearch: (<http://westminsterresearch.wmin.ac.uk/>).

In case of abuse or copyright appearing without permission e-mail repository@westminster.ac.uk

1		
2		
3		
4	Effects of upper-body, lower-body, or combined resistance training on the ratio of follistatin and myostatin	1
5	in middle-aged men	2
6		
7		
8		3
9		
10		4
11		
12	Reza Bagheri ¹ , Amir Rashidlamir ¹ , Mohamad S. Motevalli ¹ , Bradley T. Elliott ² , Javad Mehrabani ³ , Alexei Wong ⁴	5
13		
14		6
15		
16	Short title: Follistatin and Myostatin changes through resistance training	7
17		
18		8
19		
20		9
21		
22		
23	¹ Department of Exercise Physiology, Ferdowsi University of Mashhad, Mashhad, Iran.	10
24		
25		
26	² Translational Physiology Research Group, School of Life Sciences, University of Westminster, London, UK	11
27		
28	³ Department of Exercise Physiology, University of Guilan, Rasht, Iran	12
29		
30	⁴ Department of Health and Human Performance, Marymount University, Arlington, Virginia, USA	13
31		
32		14
33		
34		
35	Corresponding Author: Amir Rashidlamir, PhD; Department of Exercise Physiology, Ferdowsi University of	15
36	Mashhad, Mashhad, Iran. E-mail: Rashidlamir@um.ac.ir; Tel.: +989151514174	16
37		
38		17
39		
40		
41	COUNTING:	18
42		
43	Abstract - 249	19
44		
45		
46	Main Text - 3492	20
47		
48		21
49		
50		22
51		
52		23
53		
54		24
55		
56		25
57		
58		
59		
60		
61	1	
62		
63		
64		
65		

1
2
3
4
5
6
7 **Abstract** 1
8

9 **Purpose:** Due to the mechanistic role of myostatin and follistatin in modulating muscle mass, shifts in the 3
10 follistatin to myostatin ratio (F: M) may help explain changes in muscular size in response to resistance training 4
11 (RT). The present study examined whether differential responses in follistatin and myostatin occur based on the 5
12 amount of active musculature in a RT program in middle-aged men. 6

13
14
15 **Methods:** Forty middle-aged men (age= 46.5±3.1 years) were randomly assigned to one of 4 groups, upper-body 7
16 RT (UB; n=10), lower-body RT (LB; n=10), combined RT (UB+LB; n=10) or control (C; n=10). The training 8
17 protocol consisted of 3 exercise sessions per week for 8 weeks. Blood samples were obtained at baseline and 48 9
18 hours after the final session of the training program. 10

21
22 **Results:** Muscle mass significantly increased (p<0.05) following UB= 0.76 ± 0.46 kg, LB= 0.90 ± 0.29 kg, 11
23 UB+LB= 1.38 ± 0.70 kg, compared to no changes after control. Serum follistatin increased in the LB= 0.24 ± 0.06 12
24 ng.mL⁻¹, UB= 0.27 ± 0.17 ng.mL⁻¹, UB+LB= 0.50 ± 0.18 ng.mL⁻¹, while serum myostatin decreased in the LB= - 13
25 0.11 ± 0.08 ng.mL⁻¹ and UB+LB= -0.34 ± 0.23 ng.mL⁻¹, but not UB= 0.07 ± 0.16 ng.mL⁻¹. Further, change in 14
26 concentration following training was larger between UB+LB and either LB or UB alone for both follistatin and 15
27 myostatin. 16

28
29
30
31
32 **Conclusions:** Both UB and LB increase muscle mass and alter the F: M ratio, however the change in these 17
33 endocrine markers is approximately twice as large if UB and LB is combined. The endocrine response to RT of 18
34 myostatin and follistatin may depend on the volume of muscle mass activated during training. 19

35
36
37 **Keywords:** Follistatin, Myostatin, Resistance training, Follistatin to Myostatin ratio. 20

38
39 **Abbreviations:** 21

40
41 RT= Resistance training 22

42
43 F: M= Follistatin to myostatin ratio 23

44
45 SkMM= Skeletal muscle mass 24

46
47 PBF= Percent body fat 25

48
49 BMI=Body mass index 26

50
51 LBM=Lean body mass 27

52
53 FM=Fat mass 28

Introduction

Increasing age is associated with gradual, progressive and spontaneous erosive changes in most physiological systems and functions of the body. It has been well demonstrated that a substantial decrease in muscle strength occurs with aging, which decreases the ability to perform daily activities in middle-age and older adults (Viitasalo et al. 1985). To combat these decrements in muscular performance and functional ability, it is recommended that middle-aged and older adults perform regular resistance training (RT) (Hakkinen et al. 2000). It is well established that RT is an effective modality for increasing muscular strength, hypertrophy, and power (Frontera et al. 1988). Based on available evidence, regular RT in middle-aged adults induces muscle hypertrophy, which positively influences muscular strength and quality of life, both in the immediate period, and ideally into older age (Law et al. 2016). Recently, the influence of RT on myokines and adipo-myokines involved in the stimulation and inhibition of the muscular hypertrophic response has been explored. One notable myokine is myostatin, a powerful negative regulator of muscle size (Elliott et al. 2012). Myostatin circulates in the bloodstream, binds to Activin Type II receptors in the muscle, thereby instigating intercellular signaling pathways that inhibit muscle growth (Gonzalez-Cadavid et al. 1998). Whilst a logical hypothesis would be for an increase in serum myostatin with advancing age, there is currently mixed evidence for this. Indeed, while early work suggested that serum myostatin levels increase with advancing age (Yarasheski et al. 2002), more recent examinations have suggested no difference in circulating myostatin concentration in younger vs older men with and without sarcopenia (Ratkevicius et al. 2011) nor in a cross-sectional study of individuals spanning 18-68 years of age (Barrios-Silva et al. 2018). RT can reduce the expression of myostatin at the mRNA or protein level (Hulmi et al. 2007), leading to exercise-induced muscle hypertrophy in the healthy humans. In fact, many studies have shown that RT can generate a significant decrease in myostatin levels (Allen et al. 2011; Laurentino et al. 2012). For instance, It has revealed that serum myostatin levels decreased 10% by enzyme-linked immunosorbent assay (ELISA) after 12-week arm and leg press RT (Saremi et al. 2010). Also, It has been reported reduced myostatin mRNA expression in young and old men and women in response to 9 weeks of RT (Roth et al. 2003). The mechanism responsible for the RT-induced blunting of myostatin may involve the myostatin-inhibitor follistatin. This glycoprotein, blocks the myostatin receptor, thus decreases myostatin's effect and increasing muscle mass (Tortoriello et al. 2001). In the presence of follistatin, myostatin is unable to bind to its own receptor, and its atrophic actions are inhibited. In a study, the authors demonstrated that 12-week high-intensity RT increased serum follistatin-like related gene levels, a homolog of follistatin in certain physiological mechanisms, including myostatin inhibition (Willoughby 2004). Thus, the ratio of follistatin to myostatin (F:M) has become a commonly reported important factor in studies examining body composition (i.e. lean mass and fat mass) and the alterations of muscle strength that occur with chronic training (Rodgers and Garikipati 2008; Tortoriello et al. 2001). It was investigated the effect of eight weeks of two different intensities of RT (low or high) on F: M ratio in 21 sedentary young women. They showed that F:M ratio increased dramatically in high-intensity group (Attarzadeh Hosseini et al. 2017). Additionally, it has observed an increase of F:M ratio after 8 weeks of blood flow restriction (KAATSU) training in active men (Laurentino et al. 2012). However, although several studies report an increase in the F:M ratio with RT, such changes have not been found

1
2
3
4 in all investigations (de Souza et al. 2014; Schiffer et al. 2011). This discrepancy might be related to different 1
5 variables within a RT program, including the amount of active musculature involved. It is likely that the total 2
6 amount of active musculature plays a role in adaptations to RT, and current recommendations advocate the training 3
7 of the entire body's musculature (Medicine 2009). However, limited information is available concerning the 4
8 impact of the quantity of active musculature on the F: M ratio. Relative to reports in younger adults, less work has 5
9 examined the effect of training on myostatin and follistatin in older cohorts. **The Vienna Active Ageing cohort** 6
10 **reported no change in myostatin following 6 months of RT in a cohort of older women (65 – 92 years of age). RT** 7
11 **was performed twice a week and consisted of various elastic band exercises for the major muscle groups. Intensity** 8
12 **was progressed by increasing the resistance of elastic band.** (Hofmann et al. 2016). **Whilst** chronically trained 9
13 masters athletes show similar concentrations of circulating serum myostatin as age matched untrained controls 10
14 (Elliott et al. 2017) . As myostatin appears unresponsive to acute and chronic training in an older cohort, early and 11
15 sustained (e.g. young-adult to middle aged) RT interventions may be needed for functional maintenance of muscle 12
16 mass and strength during ageing. Therefore, the purpose of the present investigation was to determine the 13
17 responses in follistatin and myostatin following RT of differing muscle volumes in middle-aged men. It was 14
18 hypothesized that the RT program involving the greatest quantity of muscle mass (i.e. full-body training) would 15
19 elicit greater increases in follistatin and decreases in myostatin as compared to programs involving only select 16
20 portions of the body's musculature (i.e. upper body or lower body training only). 17
21
22
23
24
25
26
27
28
29
30
31
32

33 **Methods** 19

34 **Participants** 20

35
36
37 Forty middle-aged men (40-53 years) participated on the present study. Exclusion criteria included cardiovascular 21
38 diseases, diabetes, hypertension or other risk factors based on a doctor's examination. All men were sedentary, and 22
39 self-reported performing less than 1 hour of exercise per week in the previous year. **Participants were not taking** 23
40 **any supplements or medications, including non-steroidal anti-inflammatory drugs.** All participants gave written 24
41 informed consent before their inclusion in the study. The study protocol was approved by the Institutional Human 25
42 Subject committee and carried out in accordance with the Declaration of Helsinki. 26
43
44
45
46
47
48
49

50 **Study Design** 28

51 Before baseline measurements, participants were familiarized with the study tests and procedures and were 29
52 randomly assigned to into one of four groups: upper body RT (UB; n=10), lower body RT (LB; n=10), combined 30
53 upper and lower body RT (UB+LB; n=10) or control (C; n=10) groups. **Allocation was stratified by BMI (<25.0 or** 31
54 **≥25.0 kg/m²), and the sequence was randomized by a computer.** Measurements were collected at baseline and after 32
55 8 weeks during the same time of day (-1 hour). Participants were instructed not to alter their regular lifestyle and 33
56 dietary habits during the study. 34
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6 Anthropometry and Body Composition 1
2

7
8 Upon entering the laboratory, participants were asked to urinate (void) completely within 30 min of the test and 3
9 then had their body weight measured with a digital scale (lumbar, China) to the nearest 0.1 kg. Participant's height 4
10 was measured with a stadiometer (Race industrialization, China) to the nearest 0.1 cm. 5

11
12
13 BMI, PBF, SkMM and LBM were evaluated by multi-frequency bioelectrical impedance device (Inbody 720, 6
14 South Korea; Table 1). We instructed the participants to fast for 12 hours (an overnight fast, with at least 8 hours of 7
15 sleep) and refrain from physical activity for the previous 36 hours before the test. The participants were also 8
16 instructed to avoid exercising consuming alcohol for 48 hours before the test. 9
17
18
19
20
21
22

23 Blood Sampling and Analysis 11

24 Fasting blood samples (5 mL) were obtained from the cubital vein using standard procedures. The initial collection 12
25 occurred 48 hours before the baseline training session. After clotting occurred, blood samples were centrifuged at 13
26 3000 RPM for 10 minutes. Spun serum was removed from the centrifuge and frozen at -70°C for later analysis. 14
27 Serum myostatin (bound) (human myostatin, Glory Science Co, Del Rio, TX, USA) and follistatin (human 15
28 follistatin, Glory Science Co, Del Rio, TX, USA) concentrations were measured using enzyme-linked 16
29 immunosorbent assay (ELISA) kits according to manufacturer instructions. The intraassay and interassay coefficient 17
30 of variation were 8.1% and 4.5%, 8.5% and 5.4% for myostatin and follistatin, respectively. 18
31
32
33
34
35
36
37

38 Strength Testing 20

39 1RM testing 21

40
41 Maximal strength testing took place 24 hours after the body composition measurement. 1RM testing was 22
42 performed to determine training intensity for our resistance training protocol. Before the beginning of the test, 23
43 authors explained all the purpose, attendant risks, and discomforts, responsibilities of the participant, benefits, 24
44 inquiries and freedom of consent for all of them in Ferdowsi university of Mashhad. They were instructed to refrain 25
45 from alcohol for 48 hours, caffeinated drinks for 12 hours and food intake for 2 hours before the testing session; 26
46 however, water consumption was allowed. Participants warmed up their bodies for 10 minutes of general (5 27
47 minutes slow running on treadmill; 3 to 5 km speed, or elliptical; with 5 to 10 level) and specific warm-up 28
48 activities (5 minutes, e.g. medicine ball twist 1x10, medicine ball wood chops 1x10, straddled toe touch 2x5, 29
49 Dynamic quadriceps stretch 1 x 5, Medicine ball squat 1 x 5 to 8) before the test. The participants performed 2 30
50 attempts and their highest lifted weight and number of repetitions was recorded. The number of repetitions to 31
51 fatigue did not exceed 10. There was a 3 to 5 minutes between attempts. After the testing session, the participant's 32
52 maximal strength was predicted using the following formula: 33
53
54
55
56
57
58
59
60
61
62
63
64
65

1RM= weight/ (1.0278-0.0278×reps) (Murach and Bagley 2016):

Isometric quadriceps strength test

The participants stood on the body of the device and bent the knees to 130 to 140 degrees and held the trunk upright. The participant grasped the dynamometer handle with the pronation position and adjusted the length of the chain so that the handle of the dynamometer was placed on the thigh. The existing belt attached to the participants shoulders and attached it to each side of the handle of the dynamometer. This allowed the direct quantification of quadriceps isometric force. The participants performed the test 3 times each time and rested for 3 to 5 minutes between each attempt to reduce measurement error due to fatigue. All three attempts were recorded, while the highest was used for the analysis.

Isometric handgrip strength test

The participants held the dynamometer in the hand to be tested, with the arm at right angles and the elbow by the side of the body. The handle of the dynamometer was adjusted as required, with the base resting on first metacarpal (heel of palm), while the handle rested on the middle phalange of the four fingers. When ready the participant contracted the dynamometer with maximum isometric effort, which was maintained for approximately 5 seconds. No other body movement was allowed. The participant was strongly encouraged to give a maximum effort. All three attempts were recorded, while the highest was used for the analysis (Roberts et al. 2011).

Resistance Training Protocol

Preparatory phase:

All the participants performed 1 week of RT, consisting of 3 exercise sessions, for familiarization before the main training intervention. This phase allowed for supervised instruction of proper lifting technique, familiarization with all exercises, and ensured that the participants initiated the study with a comparable training base. The adaptation phase included a total of 6 exercises (Table 3). The preparatory phase program was adapted from previous literature in middle-age men (Church et al. 2016).

Training Phase

Following the preparatory phase, participants in the UB, LB and UB+LB completed the supervised training 3 times a week, separated by at least 48 hours for 8 weeks. Before each training session, the first 10 minutes included general and specific warm-up activities (slow running, stretching and light RT). After the general warm-up, participants completed a specific warm-up of 2 sets of 20 repetitions with 30% of 1RM with a 30 seconds between sets. Following the specific warm-up, training included 3 sets per exercise in weeks 1-4 and 4 sets per exercise in weeks 5-8. The rest interval between sets was 30 seconds in the first and second week, 60 seconds in the third and fourth week, 75 seconds in fifth and sixth weeks, and 90 seconds in seventh and eighth weeks (table 4). They rest interval between exercises was 2 minutes throughout the training period.

1
2
3
4 In each RT program, all exercises included 3 sets with the intensity of 50% to 80% 1RM (Table 4). To verify the 1
5 principle of overload, the following formula was used to predict the 1RM and then determine the exercise load 2
6 before the start of the training period, the fourth week, and at the end of the eighth week: 3
7

$$1RM = W / [1.0278(0.0278.r)] \text{ (Murach and Bagley 2016)}$$

8
9
10
11

12 The exercises (in order) included in the UB training program were lateral pulldown, chest press, barbell shoulder 6
13 press, lateral raise, standing barbell biceps curl, cable triceps pushdown. The exercises included in the LB training 7
14 program were barbell squat, hack squat, lunges, leg extension, lying leg curls and standing calf raises. The 8
15 exercises included in the UB+LB program were lateral pulldown, barbell squat, chest press, lunges, lateral raise 9
16 and standing calf raise. All training sessions were completed under cautious supervision from certified trainers and 10
17 researchers. The periodized RT programs were adapted from previous literature (Simão et al. 2013), following 11
18 recommendations by the National Strength and Conditioning Association (Haff and Triplett 2015). 12
19
20
21
22
23
24

25 Nutrient intake and dietary analysis 14

26 Participants were instructed not to alter their dietary habits during the study. To minimize dietary variability, the 15
27 participants were required to submit 3-day (2 weekdays and 1 weekend) food records at baseline and at 8 weeks of 16
28 the assigned intervention. Each dietary item was entered into the program Diet Analysis Plus version 10 (Cengage, 17
29 Boston, MA, USA) and total energy consumption, and the amount of energy derived from proteins, fats, and 18
30 carbohydrates was assessed, with paired sample t-tests used to confirm a lack of change in dietary factors at pre- 19
31 and post-time points (table 5). 20
32
33
34
35
36
37
38
39

40 Statistical analysis 23

41 Normality was confirmed by the Shapiro–Wilk test. Based on previous data (Negaresh et al. 2017; Saremi et al. 24
42 2010). It was calculated that 10 participants per group would provide 80% power (2-sided $\alpha=0.05$) to detect 9% 25
43 and 7% changes in myostatin and follistatin, respectively. Differences between phenotypic measures, strength 26
44 measures and endocrine markers were compared by a 4×2 ANOVA with repeated measures (group (control x LB x 27
45 UB x UB+UB) x time (pre x post)). One-way ANOVA was used for ‘change in’ (Δ) scores between times (pre, 28
46 post) where appropriate. Post hoc testing was performed by Bonferroni corrected t-tests. SPSS (Version 24, IBM) 29
47 was used for all statistical analysis, and all figures were prepared in GraphPad Prism (Version 5.03, GraphPad 30
48 Software). 31
49
50
51
52
53
54
55

56 Results 33

57 RT alters phenotype in a training model independent manner 34
58
59
60
61
62
63
64
65

1
2
3
4 Baseline parameters between the two groups were not significantly different ($p > 0.05$). A significant group x time 1
5 interaction was noted for SkMM ($p < 0.001$) and for FM ($p < 0.001$; Figure 1A & C, respectively). SkMM had a 2
6 significantly greater increase [UB= 0.76 kg (95% confidence interval {CI}, 0.48 to 1.05); LB= 0.90 kg (95% CI, 3
7 0.72 to 1.08) and UB+LB= 1.38 kg (95% CI, 0.95 to 1.81)] and FM a decreased [UB= -1.61 kg (95% CI, -1.91 to - 4
8 1.29); LB= -1.32 kg (95% CI, -1.57 to -1.07) and UB+LB= -2.26 kg (95% CI, -3.31, to -1.21)] post-training than 5
9 did the control ($p < 0.05$ in training groups, $p > 0.05$ control group). However, no difference in absolute mass of 6
10 SkMM gained or FM lost was noted between UB+LB and either UB or LB alone ($p > 0.05$; Figure 1B & D). 7
11
12
13
14
15
16
17

18 Whole-body RT alters endocrine markers to a greater degree than upper or lower body RT alone 9

19
20 A significant group x time interaction was noted for serum concentration of myostatin, follistatin and the 10
21 myostatin:follistatin concentration ratio (myostatin $p < 0.001$, follistatin $p < 0.001$, myostatin: follistatin ratio $p <$ 11
22 0.001). Myostatin concentration was shown to be decreased ($p < 0.05$) in LB [-0.11 ng.mL⁻¹ (95% CI, -0.16 to - 12
23 0.06)] and UB+LB [-0.34 ng.mL⁻¹ (95% CI, -0.48 to -0.19)] groups, but not UB or control ($p > 0.05$; figure 2C & 13
24 D). Whilst follistatin was increased post-training [UB= 0.27 ng.mL⁻¹ (95% CI, 0.16 to 0.38); LB= 0.24 ng.mL⁻¹ 14
25 (95% CI, 0.20 to 0.28) and UB+LB= 0.50 ng.mL⁻¹ (95% CI, 0.39 to 0.61)] relative to control groups ($p < 0.05$ in 15
26 training groups; Figure 2A), of more interest was the observation that UB+LB increased follistatin to a greater 16
27 extent than UB or LB alone ($p < 0.05$; Figure 2B). Finally, the follistatin:myostatin ratio was increased in all 17
28 training groups [UB= 0.07 (95% CI, 0.04 to 0.10); LB= 0.07 ng.mL⁻¹ (95% CI, 0.06 to 0.09) and UB+LB= 0.18 18
29 (95% CI, 0.12 to 0.24)], relative to both the control group ($p < 0.05$), and to a greater degree in UB+LB than UB or 19
30 LB alone ($p < 0.05$; Figure 2E & F). 20
31
32
33
34
35
36
37
38
39
40

41 RT improves isometric strength 22

42 A significant group x time interaction ($p < 0.001$, table 2) was noted for isometric quadriceps which significant 23
43 increased ($p < 0.05$) in all training groups [UB= 2.4 kg (95% CI, 2.2 to 2.6); LB= 4.6 kg (95% CI, 4.3 to 5.0) and 24
44 UB+LB= 3.6 kg (95% CI, 3.2 to 3.9)] compared to no changes after the control group. The LB increased isometric 25
45 quadriceps strength to a greater extent than UB+LB or LB alone ($p < 0.05$). There were also significant increases 26
46 ($p < 0.05$) in right [UB= 4.6 kg (95% CI, 4.3 to 5.1); LB= 2.4 kg (95% CI, 2.1 to 2.6) and UB+LB= 2.7 kg (95% 27
47 CI, 2.4 to 2.9)] and left [UB= 3.2 kg (95% CI, 2.8 to 3.6); LB= 1.5 kg (95% CI, 1.2 to 1.8) and UB+LB= 2.3 kg 28
48 (95% CI, 2.1 to 2.6)] isometric handgrip strength in all training groups but no changes after control. The UB 29
49 increased right and left isometric handgrip strength to a greater extent than UB+LB or LB alone ($p < 0.05$) 30
50
51
52
53
54
55
56
57
58
59

60 Discussion 33

1
2
3
4 This study aimed to identify the effect of three 8-week RT protocols, each activating different volumes of muscle 1
5 mass (LB, UB or UB+LB) on follistatin, myostatin and their ratio in middle-aged men. The key finding we report 2
6 is that the response of follistatin and myostatin to muscle mass building RT is proportional to the volume of muscle 3
7 mass activated. In response to each RT program (UB+LB, UB, and LB), serum follistatin levels increased 4
8 significantly, and the greatest effect was observed in the group performing both upper- and lower-body RT (i.e. 5
9 UB+LB). The F: M ratio also increased significantly in all three groups, with the greatest effect observed in 6
10 UB+LB. Myostatin decreased similarly in all training groups, although not significantly in the UB group following 7
11 training. As would be expected, all forms of RT resulted in increased muscle mass gain. Indeed, it is well described 8
12 in the literature that RT results in increases in muscle mass in most healthy individuals (Grgic et al. 2018), and that 9
13 greater muscle gain results from higher volumes of training (Schoenfeld et al. 2016), higher number of sets 10
14 completed (Krieger 2010), and indeed specificity of training response to the muscle mass utilized in training 11
15 (Taniguchi 1997; Taniguchi 1998) . The role of myostatin and follistatin in response to RT has been reported prior 12
16 to this paper, and it is noteworthy that our results here mirror those of others. In particular, Walker et al. (2004) 13
17 showed that plasma myostatin protein levels decreased by approximately 20 % in participants after 10 weeks of 14
18 twice weekly bouts of RT, in a manner similar to that shown by others (Saremi et al. 2010). Both younger and 15
19 older men show decreased circulating myostatin and increased follistatin concentration in response to 8 week of 16
20 RT, in a highly similar manner to the results we report here (Negareh et al. 2017). However, it is noteworthy, that 17
21 when challenged with UB+LB, our participants show an endocrine response of myostatin and follistatin that is 18
22 approximately twice what is seen if UB or LB alone is performed. This novel finding suggests that the volume of 19
23 muscle mass activated is a key factor in the endocrine portion of the anabolic response to RT. Indeed, RT studies 20
24 that have examined volume of training report lower volumes produce a lesser endocrine response of anabolic 21
25 hormones such as growth hormone and testosterone (Goto et al. 2004; Gotshalk et al. 1997) . In contrast to these 22
26 results and the findings we report here, Walker et al, (2004) did not show a difference in plasma myostatin 23
27 response between a whole body vs a UB training protocol similar to our own, when semi-quantitatively examined 24
28 by Western blot. We would speculate that this may be due to the measurement technique used, and that current 25
29 ELISA based protocols would be more sensitive to changes in concentration than Western blot approaches; 26
30 however, this has not be objectively tested by ourselves to date. Alternatively, it was examined a relatively light 27
31 training workload (twice weekly) and muscle mass gain relatively to the load used in our middle-aged men, which 28
32 may underlie the differences in myostatin response by muscle volume trained (UB vs LB vs LB+UB)(Walker et al. 29
33 2004). The single-chain polypeptide follistatin is a member of the larger family of the Transforming growth factors 30
34 (Görgens et al. 2013), that is ubiquitously expressed in all tissues of the human body, including skeletal muscle, 31
35 and has paracrine and autocrine effects. Previous reports showed that follistatin has anabolic and catabolic effects 32
36 in skeletal muscle and adipose tissue, respectively (Gilson et al. 2009; Rodino- Klapac et al. 2009), likely via 33
37 follistatin's ability to bind and inactivate myostatin (Amthor et al. 2004). Most studies examining RT and 34
38 follistatin have found increased circulating follistatin concentrations in their research, which is in line with the 35
39 results we report here (Hofmann et al. 2016; Jang et al. 2016; Negareh et al. 2017). 36
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4 The increase in F: M ratio after our UB+LB as related to a change in the balanced disturbance of the 1
5 muscle growth regulators towards positive regulators. In normal conditions, to maintain muscle fiber size, there is 2
6 a homeostatic balance between positive (such as follistatin and IGF-1) and negative (such as myostatin) regulators 3
7 of muscle growth; but this balance, when the muscle is atrophic, leads to a dominance in negative regulators; 4
8 hence, loading the muscle through RT, may lead to the dominance positive regulators. Although the mechanism of 5
9 communication between these regulators is not completely clear, this connection seems to be made through a very 6
10 complex negative feedback loop (Gonzalez-Cadavid et al. 1998). Furthermore, has been shown that the release rate 7
11 of myokines and regulatory factors for muscle growth is largely dependent on the intensity and volume of muscles 8
12 involved in the activity (Motevalli et al. 2015). It appears that the volume of muscle activated is a key variable in 9
13 RT, as this promotes anabolic pathways in a larger quantity of tissue (Burd et al. 2010; Terzis et al. 2010). 10
14 Accordingly, the combined exercise protocol of this study, compared with the upper and lower trunk exercise 11
15 protocols, showed an increase in the F: M ratio in middle-aged men due to the involvement of more muscles in RT. 12
16 However, at least within the time frames reported here, this did not result in a significant difference in the gains in 13
17 whole body muscle mass or decrease in fat mass between training groups. This may be at least partially explained 14
18 by the short duration of the training protocol. It is generally held that neural adaptations predominate first, while 15
19 muscle hypertrophy comes to being evident after approximately 8 weeks of training (Haff and Triplett 2015). 16
20 Therefore, our 8-week intervention might not have warranted enough time for a significant difference in muscle 17
21 mass gains between training groups (although there was a non-significant tendency for the UB+LB to have higher 18
22 improvements than the other groups). Additionally, somewhat similar training volume was used in all 3 groups [6 19
23 exercises (3 for each the upper and lower half) in the UB+LB compared to 6 exercises for the UB and LB groups], 20
24 which potentially affected the myostatin and follistatin response and hence the muscle mass gain and fat loss in the 21
25 UB+LB. Myostatin can both inhibit or promote adipogenesis (Deng et al. 2017) which may create some confusion 22
26 as far as its effects on fat loss in our study. Nonetheless, previous results in rats show that the reduction in fat mass 23
27 in myostatin deficient rats is influenced by metabolic changes in skeletal muscle (Guo et al. 2009). Therefore, the 24
28 changes in fat mass in the UB+LB that tended to be greater (although non-significant) than the LB and UB, might 25
29 be influenced by greater improvement in muscle mass in the UB+LB group. However, future RT studies in 26
30 different populations are needed to validate our claim. 27
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45

46 The interesting findings of our investigation should be understood in the context of the following 28
47 limitations. We did not measure training motivation and fiber type, which would have strengthened the matching of 29
48 participants and hence our study design. We were not able to document training data, such as loads and changes in 30
49 work volume, that might have helped explaining some of our findings. This study is also limited by short duration 31
50 and low-moderate intensity (50-80% of 1RM) of our intervention; which warrants further investigations with longer 32
51 interventions and heavier loads with a focus on understanding the relationship between changes in SkMM and FM, 33
52 and myostatin as well as follistatin responses after combined upper- and lower-body RT. 34
53
54
55
56
57

58 In conclusion, the findings of this study show that the use of combined upper- and lower-body RT 36
59 increases the serum ratio of F: M in middle-aged men over upper or lower-body training along, suggesting that 37
60
61
62
63
64
65

1		
2		
3		
4	volume of muscle mass activated may influence the magnitude of the serum myostatin and follistatin response.	1
5	Future research might examine the volume of muscle activated vs the total amount of work completed, as this	2
6	research suggests one of these two variables underlies the endocrine anabolic response of follistatin and myostatin.	3
7		
8		
9		
10		4
11		
12	Acknowledgements	5
13		
14	The authors wish to thank all the participants in this research project.	6
15		
16		7
17		
18	Conflict of interest	8
19		
20	The authors declare no conflict of interest.	9
21		
22		
23		10
24		
25		
26		11
27		
28		12
29		
30		13
31		
32		14
33		
34		15
35		
36		16
37		
38		17
39		
40		18
41		
42		19
43		
44		20
45		
46		21
47		
48		22
49		
50		23
51		
52		24
53		
54		25
55		
56		
57	References	25
58		
59		
60		
61		
62		
63		
64		
65		

1
2
3
4 Allen DL, Hittel DS, McPherron AC (2011) Expression and function of myostatin in obesity, diabetes, and 1
exercise adaptation *Medicine and science in sports and exercise* 43:1828 2
5
6 Amthor H, Nicholas G, McKinnell I, Kemp CF, Sharma M, Kambadur R, Patel K (2004) Follistatin 3
complexes Myostatin and antagonises Myostatin-mediated inhibition of myogenesis 4
7
8 *Developmental biology* 270:19-30 5
9
10 Attarzadeh Hosseini SR, Moeinnia N, Motahari Rad M (2017) The effect of two intensities resistance 6
11 training on muscle growth regulatory myokines in sedentary young women *Obesity Medicine* 2 7
12
13 Barrios-Silva LV et al. (2018) Activin subfamily peptides predict chronological age in humans 8
14 *Physiological reports* 6:e13823 9
15
16 Burd NA et al. (2010) Resistance exercise volume affects myofibrillar protein synthesis and anabolic 10
17 signalling molecule phosphorylation in young men *The Journal of physiology* 588:3119-3130 11
18
19 Church DD et al. (2016) Comparison of high-intensity vs. high-volume resistance training on the BDNF 12
20 response to exercise *Journal of Applied Physiology* 121:123-128 13
21
22 de Souza EO et al. (2014) Effects of concurrent strength and endurance training on genes related to 14
23 myostatin signaling pathway and muscle fiber responses *The Journal of Strength & Conditioning* 15
24 *Research* 28:3215-3223 16
25
26 Deng B et al. (2017) The function of myostatin in the regulation of fat mass in mammals *Nutrition &* 17
27 *metabolism* 14:29 18
28
29 Elliott B, Renshaw D, Getting S, Mackenzie R (2012) The central role of myostatin in skeletal muscle and 19
30 whole body homeostasis *Acta Physiologica* 205:324-340 20
31
32 Elliott BT, Herbert P, Sculthorpe N, Grace FM, Stratton D, Hayes LD (2017) Lifelong exercise, but not 21
33 short-term high-intensity interval training, increases GDF11, a marker of successful aging: a 22
34 preliminary investigation *Physiological reports* 5 23
35
36 Frontera WR, Meredith CN, O'Reilly KP, Knuttgen HG, Evans WJ (1988) Strength conditioning in older 24
37 men: skeletal muscle hypertrophy and improved function *Journal of applied physiology* 64:1038- 25
38 1044 26
39
40 Gilson H, Schakman O, Kalista S, Lause P, Tsuchida K, Thissen J-P (2009) Follistatin induces muscle 27
41 hypertrophy through satellite cell proliferation and inhibition of both myostatin and activin 28
42 *American Journal of Physiology-Endocrinology and Metabolism* 297:E157-E164 29
43
44 Gonzalez-Cadavid NF et al. (1998) Organization of the human myostatin gene and expression in healthy 30
45 men and HIV-infected men with muscle wasting *Proceedings of the National Academy of* 31
46 *Sciences* 95:14938-14943 32
47
48 Görgens SW, Raschke S, Holven KB, Jensen J, Eckardt K, Eckel J (2013) Regulation of follistatin-like 33
49 protein 1 expression and secretion in primary human skeletal muscle cells *Archives of physiology* 34
50 *and biochemistry* 119:75-80 35
51
52 Goto K, Nagasawa M, Yanagisawa O, Kizuka T, Ishii N, Takamatsu K (2004) Muscular adaptations to 36
53 combinations of high-and low-intensity resistance exercises *Journal of Strength and* 37
54 *Conditioning Research* 18:730-737 38
55
56 Gotshalk LA et al. (1997) Hormonal responses of multiset versus single-set heavy-resistance exercise 39
57 protocols *Canadian Journal of Applied Physiology* 22:244-255 40
58
59 Grgic J, Mcllvenna LC, Fyfe JJ, Sabol F, Bishop DJ, Schoenfeld BJ, Pedisic Z (2018) Does Aerobic Training 41
60 Promote the Same Skeletal Muscle Hypertrophy as Resistance Training? A Systematic Review 42
61 and Meta-Analysis *Sports Medicine*:1-22 43
62
63 Guo T, Jou W, Chanturiya T, Portas J, Gavriloova O, McPherron AC (2009) Myostatin inhibition in muscle, 44
64 but not adipose tissue, decreases fat mass and improves insulin sensitivity *PloS one* 4:e4937 45
65
66 Haff GG, Triplett NT (2015) *Essentials of strength training and conditioning* 4th edition. Human kinetics, 46

1
2
3
4 Hakkinen K, Pakarinen A, Kraemer WJ, Newton RU, Alen M (2000) Basal concentrations and acute 1
5 responses of serum hormones and strength development during heavy resistance training in 2
6 middle-aged and elderly men and women *Journals of Gerontology-Biological Sciences and* 3
7 *Medical Sciences* 55:B95 4
8
9 Hofmann M et al. (2016) Effects of elastic band resistance training and nutritional supplementation on 5
10 muscle quality and circulating muscle growth and degradation factors of institutionalized elderly 6
11 women: the Vienna Active Ageing Study (VAAS) *European journal of applied physiology* 116:885- 7
12 897 8
13
14 Hulmi JJ et al. (2007) Postexercise myostatin and activin IIb mRNA levels: effects of strength training 9
15 *Medicine & Science in Sports & Exercise* 39:289-297 10
16
17 Jang KS, Kang S, Woo SH, Bae JY, Shin KO (2016) Effects of combined open kinetic chain and closed 11
18 kinetic chain training using pulley exercise machines on muscle strength and angiogenesis 12
19 factors *Journal of physical therapy science* 28:960-966 13
20
21 Krieger JW (2010) Single vs. multiple sets of resistance exercise for muscle hypertrophy: a meta-analysis 14
22 *The Journal of Strength & Conditioning Research* 24:1150-1159 15
23
24 Laurentino GC et al. (2012) Strength training with blood flow restriction diminishes myostatin gene 16
25 expression *Med Sci Sports Exerc* 44:406-412 17
26
27 Law TD, Clark LA, Clark BC (2016) Resistance exercise to prevent and manage sarcopenia and dynapenia 18
28 *Annual review of gerontology & geriatrics* 36:205 19
29
30 Medicine ACoS (2009) American College of Sports Medicine position stand. Progression models in 20
31 resistance training for healthy adults *Medicine and science in sports and exercise* 41:687 21
32
33 Motevalli MS, Dalbo VJ, Attarzadeh RS, Rashidlamir A, Tucker PS, Scanlan AT (2015) The effect of rate of 22
34 weight reduction on serum myostatin and follistatin concentrations in competitive wrestlers 23
35 *International journal of sports physiology and performance* 10:139-146 24
36
37 Murach KA, Bagley JR (2016) Skeletal muscle hypertrophy with concurrent exercise training: contrary 25
38 evidence for an interference effect *Sports medicine* 46:1029-1039 26
39
40 Negaresh R, Ranjbar R, Habibi A, Mokhtarzade M, Fokin A, Gharibvand M (2017) The effect of resistance 27
41 training on quadriceps muscle volume and some growth factors in elderly and young men 28
42 *Advances in gerontology= Uspekhi gerontologii* 30:880-887 29
43
44 Ratkevicius A et al. (2011) Serum concentrations of myostatin and myostatin-interacting proteins do not 30
45 differ between young and sarcopenic elderly men *Journals of Gerontology Series A: Biomedical* 31
46 *Sciences and Medical Sciences* 66:620-626 32
47
48 Roberts HC, Denison HJ, Martin HJ, Patel HP, Syddall H, Cooper C, Sayer AA (2011) A review of the 33
49 measurement of grip strength in clinical and epidemiological studies: towards a standardised 34
50 approach *Age and ageing* 40:423-429 35
51
52 Rodgers BD, Garikipati DK (2008) Clinical, agricultural, and evolutionary biology of myostatin: a 36
53 comparative review *Endocrine reviews* 29:513-534 37
54
55 Rodino-Klapac LR, Haidet AM, Kota J, Handy C, Kaspar BK, Mendell JR (2009) Inhibition of myostatin with 38
56 emphasis on follistatin as a therapy for muscle disease *Muscle & Nerve: Official Journal of the* 39
57 *American Association of Electrodiagnostic Medicine* 39:283-296 40
58
59 Roth SM, Martel GF, Ferrell RE, Metter EJ, Hurley BF, Rogers MA (2003) Myostatin gene expression is 41
60 reduced in humans with heavy-resistance strength training: a brief communication *Experimental* 42
61 *biology and medicine* 228:706-709 43
62
63 Saremi A, Gharakhanloo R, Sharghi S, Gharaati M, Larijani B, Omidfar K (2010) Effects of oral creatine 44
64 and resistance training on serum myostatin and GASP-1 *Molecular and cellular endocrinology* 45
65 317:25-30 46

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Schiffer T, Geisler S, Sperlich B, Strüder H (2011) MSTN mRNA after varying exercise modalities in humans *International journal of sports medicine* 32:683-687

Schoenfeld BJ, Ogborn D, Krieger JW (2016) Effects of resistance training frequency on measures of muscle hypertrophy: a systematic review and meta-analysis *Sports Medicine* 46:1689-1697

Simão R et al. (2013) Influence of upper-body exercise order on hormonal responses in trained men *Applied Physiology, Nutrition, and Metabolism* 38:177-181

Taniguchi Y (1997) Lateral specificity in resistance training: the effect of bilateral and unilateral training *European journal of applied physiology and occupational physiology* 75:144-150

Taniguchi Y (1998) Relationship between the modifications of bilateral deficit in upper and lower limbs by resistance training in humans *European journal of applied physiology and occupational physiology* 78:226-230

Terzis G, Spengos K, Mascher H, Georgiadis G, Manta P, Blomstrand E (2010) The degree of p70S6k and S6 phosphorylation in human skeletal muscle in response to resistance exercise depends on the training volume *European journal of applied physiology* 110:835-843

Tortoriello DV, Sidis Y, Holtzman DA, Holmes WE, Schneyer AL (2001) Human follistatin-related protein: a structural homologue of follistatin with nuclear localization *Endocrinology* 142:3426-3434

Viitasalo J, Era P, Leskinen A-L, Heikkinen E (1985) Muscular strength profiles and anthropometry in random samples of men aged 31–35, 51–55 and 71–75 years *Ergonomics* 28:1563-1574

Walker KS, Kambadur R, Sharma M, Smith HK (2004) Resistance training alters plasma myostatin but not IGF-1 in healthy men *Medicine and science in sports and exercise* 36:787-793





Willoughby DS (2004) Effects of an alleged myostatin-binding supplement and heavy resistance training on serum myostatin, muscle strength and mass, and body composition *International journal of sport nutrition and exercise metabolism* 14:461-472







Yarasheski K, Bhasin S, Sinha-Hikim I, Pak-Loduca J, Gonzalez-Cadavid N (2002) Serum myostatin-immunoreactive protein is increased in 60-92 year old women and men with muscle wasting *Journal of Nutrition Health and Aging* 6:343-348

1
2
3
4 **Figures Legends**

5 1
6 2
7
8 Figure 1: RT increases skeletal muscle mass and decreases fat mass independent of training modality. A) Skeletal 3
9 muscle mass (SkMM; kg) by group pre and post training. B) Change in (Δ) SkMM (kg) from pre to post training. C) 4
10 Fat mass (FM; kg) by group pre and post training. D) Δ FM (kg) from pre to post training. Error bars represent 5
11 standard deviation. Individual data points as shown. * indicates differences from groups as shown, ^ indicates 6
12 differences from control. Upper body (UB), lower body (LB), upper body and lower body combined (UB+LB). 7

13
14 8
15
16 Figure 2: RT alters endocrine factors myostatin and follistatin as a function of the volume of muscle mass trained. 9
17 A) Follistatin ($\text{ng}\cdot\text{mL}^{-1}$). B) Change in (Δ) follistatin pre to post ($\text{ng}\cdot\text{mL}^{-1}$). C) Myostatin ($\text{ng}\cdot\text{mL}^{-1}$). D) Δ myostatin 10
18 ($\text{ng}\cdot\text{mL}^{-1}$). E) follistatin:myostatin (F:M) ratio. F) Δ F: M ($\text{ng}\cdot\text{mL}^{-1}$). Error bars represent standard deviation. 11
19 Individual data points as shown. * indicates differences from groups as shown, ^ indicates differences from control. 12
20 Upper body (UB), lower body (LB), upper body and lower body combined (UB+LB). 13





Table

Week	Intensity (1RM)	Repetitions	Sets	Rest (second)
1	50%	15	3	30
2	50%	15	3	30
3	60%	12	3	60
4	60%	12	3	60
5	70%	10	4	75
6	70%	10	4	75
7	80%	8	4	90
8	80%	8	4	90

Table 4. Variables in the resistance training protocols

Table

Variable	Group	Pre-training	Post-training	P-value
		M±SD	M±SD	
Age (year)	UB+LB	45.6 ± 3.2	---	---
	UB	47.9 ± 1.9	---	---
	LB	45.3 ± 2.8	---	---
	C	47.5 ± 3.9	---	---
Height (cm)	UB+LB	175.9 ± 4.2	---	---
	UB	177.8 ± 2.5	---	---
	LB	174.3 ± 3.9	---	---
	C	176.7 ± 2.7	---	---
Weight (Kg)	UB+LB	83.7 ± 6.1	82.2 ± 5.9	0.00
	UB	86.8 ± 5.0	85.9 ± 4.8	0.004
	LB	81.1 ± 5.3	80.1 ± 5.3	0.001
	C	85.4 ± 5.5	85.8 ± 5.6	0.23
BMI (kg/m ²)	UB+LB	27.1 ± 1.1	26.6 ± 1.1	0.00
	UB	27.5 ± 1.6	27.2 ± 1.6	0.004
	LB	26.7 ± 1.4	26.4 ± 1.5	0.001
	C	27.4 ± 1.5	27.5 ± 1.5	0.25
LBM (kg)	UB+LB	63.2 ± 3.9	64.1 ± 3.9	0.01
	UB	64.1 ± 3.7	64.4 ± 3.9	0.18
	LB	60.7 ± 4.0	61 ± 4.1	0.07
	C	64.1 ± 4.0	64.0 ± 3.9	0.92
PBF (%)	UB+LB	24.3 ± 2.5	21.8 ± 2.4	0.001
	UB	26.1 ± 1.2	25 ± 1.6	0.007
	LB	25.2 ± 1.8	23.8 ± 1.6	0.00
	C	25.2 ± 1.8	25.2 ± 2.5	0.54
SkMM (kg)	UB+LB	35.9 ± 2.7	37.4 ± 2.4	0.00
	UB	36.1 ± 1.8	36.8 ± 2.1	0.001
	LB	34.4 ± 1.8	35.3 ± 1.7	0.00
	C	35.0 ± 2.7	34.9 ± 2.6	0.57
FM (kg)	UB+LB	20.4 ± 3.0	18.1 ± 2.8	0.001
	UB	22.7 ± 1.8	21.5 ± 1.8	0.004
	LB	20.4 ± 2.1	19.1 ± 1.9	0.00
	C	21.4 ± 2.3	21.7 ± 2.9	0.41

Table 1. Physiological characteristics of the participants. Abbreviations: BMI, Body Mass Index; LBM, Lean Body Mass; PBF, Percent Body Fat; SkMM, Skeletal Muscle Mass; FM, Fat Mass.

Table

	Group	Pre-training	Post-training	P-value
Quadriceps strength test (kg)	UB+LB	66.1 + 5.1	69.8 + 4.2	0.00
	UB	67.4 + 6.4	69.8 + 6.9	0.00
	LB	67.3 + 5.9	71.9 + 5.8	0.00
	C	69.3 + 4.8	68.5 + 5.4	0.19
Handgrip strength test (left hand; kg)	UB+LB	42.9 + 2.8	45.2 + 2.8	0.00
	UB	42.3 + 2.3	45.4 + 3.1	0.001
	LB	41.8 + 4.2	43.3 + 4.8	0.01
	C	42.3 + 3.5	42 + 4.3	0.49
Handgrip strength test (right hand; kg)	UB+LB	44.5 + 2.3	47.2 + 2.2	0.00
	UB	46.6 + 2.8	51.3 + 3.5	0.00
	LB	44.2 + 2.1	46.6 + 1.8	0.00
	C	45.8 + 2.4	45.4 + 3.4	0.47

Table 2. Values for the Quadriceps and Handgrip strenght tests

Table

Week	Exercises	Intensity (1RM)	Repetitions	Rest intervals	Sets
1	BS, LP, LE, R, SLC, LR	40	12	30 sec	3

Table 3. Adaptation phase training protocol. Abbreviations: Barbell squat, BS; Lateral raise, LR; Leg Extension, LE; Seated Leg Curl, SLC; Lat pulldown, LP; Rowing, R.

Table

	Group	Pre-training	Post-training	p-value
Energy (kcal/kg/d)	UB+LB	26.1 + 3.8	25 + 3.3	0.41
	UB	26.7 + 4.6	26 + 5.1	0.56
	LB	26.6 + 5.4	25.5 + 3.2	0.43
	C	26.2 + 5.6	26.3 + 5.7	0.97
Protein (g/kg/d)	UB+LB	0.83 + 0.13	0.86 + 0.09	0.51
	UB	0.84 + 0.18	0.85 + 0.23	0.73
	LB	0.85 + 0.18	0.85 + 0.16	0.96
	C	0.83 + 0.16	0.83 + 0.13	0.91
Fat (g/kg/d)	UB+LB	0.84 + 0.21	0.81 + 0.19	0.78
	UB	0.8 + 0.18	0.79 + 0.22	0.84
	LB	0.78 + 0.23	0.76 + 0.17	0.79
	C	0.8 + 0.29	0.8 + 0.24	0.81
CHO (g/kg/d)	UB+LB	3.8 + 0.54	3.5 + 0.3	0.25
	UB	4.0 + 0.7	3.8 + 0.8	0.42
	LB	4.0 + 0.7	3.8 + 0.6	0.34
	C	3.9 + 0.8	3.9 + 0.9	0.87

Table 5. Energy and macronutrients (Mean ± SD). Abbreviations: kcal, kilocalories; kg, kilogram of body weight; d, day; g, grams.

Author Contribution Statement

Dr. Rashidlamir conceived and designed research.

Reza Bagheri conducted the experiments and wrote the paper.

Dr. Wong and Mehrabani fixed grammatical mistakes. In addition, Dr. Wong analyzed data, contributed in writing the paper and preparing revisions.

M Motevalli analyzed the nutrition data.

Dr. Elliott analyzed the other data, wrote the results sections and fixed all the section of the paper academically.

Finally all authors read and approved the manuscript.