

Isolation of a Klebsiella bacterium-bacteriophage combination from human caecum effluent

**Lesley Hoyles¹
Anne L. McCartney²
Douwe van Sinderen³**

¹ Department of Microbiology, University College Cork, Ireland (Now Faculty of Science and Technology, University of Westminster)

² Department of Food and Nutritional Sciences, University of Reading, UK

³ Department of Microbiology, University College Cork, Ireland

This is a copy of the poster presented at the SGM Autumn Conference, University of York, UK, September 2011.

Copyright © 2011 The Authors.

The WestminsterResearch online digital archive at the University of Westminster aims to make the research output of the University available to a wider audience. Copyright and Moral Rights remain with the authors and/or copyright owners.

Whilst further distribution of specific materials from within this archive is forbidden, you may freely distribute the URL of WestminsterResearch:
(<http://westminsterresearch.wmin.ac.uk/>).

In case of abuse or copyright appearing without permission e-mail
repository@westminster.ac.uk

Isolation of a *Klebsiella* bacterium–bacteriophage combination from human caecum effluent

Lesley Hoyles¹ | Anne L. McCartney² | Douwe van Sinderen¹

Introduction

The human gastrointestinal (GI) tract harbours a diverse range of bacteria, viruses and eukaryotes, with this microbial consortium referred to as the GI microbiota. A very large body of work exists in relation to the microbiota associated with the human colon, but little is known about the microbiota associated with the caecum.

The caecum is a pouch that connects the ileum to the proximal colon, and is considered to be the beginning of the large intestine. The mucosal surface of the human caecum is heavily populated with bacteria, with substantially more biofilm formation in this region of the GI tract than the proximal or transverse colon on the basis of the examination of the colon contents of one deceased organ donor (Bollinger *et al.*, 2007, *J Theor Biol* **249**, 826–831). Colonoscopic examination of the caecum reveals the presence of highly mucoid material lining its mucosal surface (Fig. 1), from which aerobic and anaerobic bacteria can be easily isolated (unpublished data). Therefore, the biofilm associated with the caecum represents “*adherent colonies of microbes growing within an extracellular matrix*” (Bollinger *et al.*, 2007).




Fig. 1. Appearance of the caecum as viewed from the proximal colon (left panel). The yellow mucoid material lining the surface of the caecum was collected by suction during a routine colonoscopy procedure, and transported under anaerobic conditions to the University of Reading in a trap container (right panel). Ethical approval for the collection of caecum effluent from patients was obtained from St Thomas' Hospital Research Ethics Committee (06/Q0702/74) covering Guy's and St Thomas' Hospitals and transferred by agreement to London Bridge Hospital.

During a study of the microbiota associated with the caecum of patients with irritable bowel syndrome (IBS), attempts were made to isolate bacterium–bacteriophage combinations from samples of caecum effluent. Herein, we report the isolation of the first such combination from the caecum effluent of one patient.

Methods

Isolation of strain L4-FAA5 and bacteriophage Φ KLPN1 from caecum effluent. The sample was processed within 3 h of collection. Strain L4-FAA5 was isolated from diluted caecum effluent at 37 °C on fastidious anaerobe agar under anaerobic conditions. The remaining neat caecum effluent was diluted with sterile 0.5 % Lablemco/6 % NaCl, and homogenized in a stomacher for 2 min at 'high' speed, and placed on ice for 1 h. The homogenate was centrifuged at 4500 g for 20 min at 4 °C, and the supernatant passed through sterile 0.45 μ m cellulose acetate filters; the sterile filtrate contained the virus-like particles isolated from the caecum sample. Bacteria isolated were identified by determining their 16S rRNA gene sequences. *Klebsiella* strains were grown in tryptone soya medium, and used in spot-overlay assays with the filtered caecum sample and double-agar-overlay-plaque assays with the purified bacteriophage. Various characteristics of the bacteriophage were determined (Fig. 2), as was its ability to infect a panel of 20 *Klebsiella pneumoniae* subsp. *pneumoniae* clinical isolates representing various capsule types (K2, K5, K20, K54 and K57). Transmission electron microscopy was used to examine the morphology of Φ KLPN1 (Fig. 3).




Fig. 2. Characteristics of plaques of bacteriophage Φ KLPN1 when infecting strain L4-FAA5. (a) Appearance of plaques upon initial isolation from caecum effluent. (b) Increase in the size of halo surrounding plaques over the course of 4 days when cultures were incubated aerobically at 37 °C, although the central clear area of the plaques remained 2 mm in diameter. (c) Halo morphology was dependent on atmospheric conditions under which cultures were incubated. After 24 h, aerobically grown L4-FAA5 infected with Φ KLPN1 produced haloes with concentric circles (left panel), whereas anaerobically grown infected cultures produced less distinct, but uniform haloes that were larger than those produced aerobically (11 mm vs 9 mm) (right panel). The two images have been manipulated to allow the halo around the plaque in the anaerobically grown infected culture to be seen.

Results and Discussion

Five of the ten colonies isolated from the caecum sample were identified as *Klebsiella pneumoniae*, representing ~11 % of the total microbiota (2.5×10^7 of 2.2×10^8 cfu/ml caecum effluent). Strain L4-FAA5 was selected for further characterization. Using the typing method of Turton *et al.* (2010, *J Med Microbiol* **59**, 541–547), strain L4-FAA5 was identified as *K. pneumoniae* subsp. *pneumoniae* capsule type K2, *rmpA*⁺. When grown anaerobically for 24 h, strain L4-FAA5 was highly mucoid, forming amorphous colonies upon prolonged incubation. Colonies of aerobically grown cultures were 3 mm in diameter after 24 h, shiny and convex; although the colonies increased in size after prolonged incubation, they maintained their form.

Plaques identical to those shown in Fig. 2(a) were observed with all *K. pneumoniae* isolates in the spot-overlay assays with filtered caecum effluent. Strain L4-FAA5 was used as the host bacterium on which to purify and propagate the bacteriophage, Φ KLPN1. Φ KLPN1 formed clear plaques of 2 mm diameter within 3 h of inoculating an agar overlay, and was detected at $2 \times 10^5 \pm 2.65 \times 10^3$ pfu/ml caecum effluent ($n=3$). After prolonged incubation, the area around the plaques developed opaque haloes, suggesting Φ KLPN1 possesses an exopolysaccharide depolymerase (Fig. 2b). The concentric circles seen within the halo area of aerobically grown infected cultures were absent from anaerobically grown cultures (Fig. 2c). At an MOI of 0.008 and with aerobic incubation, 95 % of infectious particles adsorbed to strain L4-FAA5 (OD_{660} 0.4) within 5 min; burst size was ~2000 pfu. On the basis of its restriction profile (not shown), the genome of Φ KLPN1 was ~35 kb in size. SDS-PAGE analysis of structural proteins showed three major bands at 45, 30 and 25 kDa (not shown). Examination by transmission electron microscopy showed Φ KLPN1 to be a member of the *Siphoviridae* (Fig. 3).




Fig. 3. Transmission electron micrographs of bacteriophage Φ KLPN1. The bacteriophage has a flexible, non-contractile tail, and belongs to the *Siphoviridae*. At the terminus of the striated tail are what appear to be short fibres. The capsid is ~64 nm at its widest point, tapering to ~54 nm; the tail is ~196 nm long.

When screened against 20 clinical isolates of *K. pneumoniae* subsp. *pneumoniae*, Φ KLPN1 infected 5/6 K2 strains, but none of the other strains tested (spot-overlay assays). The ability of Φ KLPN1 to infect K2 strains suggests the bacteriophage could have therapeutic potential in nosocomial infections. Consequently, the human GI tract should be viewed as a source of bacteriophages of medical interest.

The depolymerase activity of Φ KLPN1 on strain L4-FAA5 may be of ecological significance. We have found high concentrations of bacteria within caecum biofilm material. This material may be a preferred site for bacteriophage reproduction compared to the less accessible bacteria found in luminal contents of the colon. Φ KLPN1 may be a prophage of L4-FAA5 that has reverted to a lytic cycle of infection due to the cultivation of its host outside the human body. Within the caecum, the bacteriophage may regulate population numbers within the biofilm in response to environmental changes within this ecosystem.

Acknowledgements

Jeremy D. Sanderson (Department of Gastroenterology, Guy's and St Thomas' NHS Foundation Trust and King's College London, UK) is thanked for obtaining the caecum sample. Jane F. Turton (Health Protection Agency, Colindale, UK) is thanked for her help with characterizing strain L4-FAA5 and for providing access to the clinical isolates. Horst Neve (Max Rubner – Institut, Institute of Microbiology and Biotechnology, Kiel, Germany) is thanked for providing the TEM image of bacteriophage Φ KLPN1. L.H. holds a Government of Ireland Postdoctoral Fellowship in Science, Engineering and Technology (EMPOWER scheme).

Contact information

Email: l.hoyles@ucc.ie or lesley.hoyles@reading.ac.uk